



Short Original

Non-dipping blood pressure pattern in pediatricians during on-duty

Javier Gonzalez-Garcia^{a,*}, Juan Mayordomo-Colunga^{b,f,g}, Manuel Gorostidi^{c,d}, Belen Gomez-Gonzalez^e, Eva Canteli^c, Corsino Rey^{g,h,i,j}

^a Department of Pediatrics, Hospital Universitario de Cabueñes, Gijón, Spain

^b Pediatric Intensive Care Unit, Hospital Universitario Central de Asturias, Oviedo, Spain

^c Department of Nephrology, Hospital Universitario Central de Asturias, Oviedo, Spain

^d Department of Nephrology, University of Oviedo, Oviedo, Spain

^e Department of Pediatrics, Hospital Valle del Nalón, Langreo, Spain

^f Network Biomedical Research Center, Respiratory Diseases, Institute of Health Carlos III, Madrid, Spain

^g Health Research Institute of the Principado de Asturias, Oviedo, Spain

^h Department of Pediatrics, Hospital Universitario Central de Asturias, Oviedo, Spain

ⁱ Department of Pediatrics, University of Oviedo, Oviedo, Spain

^j Primary Care Interventions to Prevent Maternal and Child Chronic Diseases of Perinatal and Developmental Origin (RICORS), Institute of Health Carlos III, Madrid, Spain

ARTICLE INFO

Article history:

Received 8 March 2022

Accepted 30 May 2022

Available online 22 June 2022

Keywords:

Hypertension

Non-dipping

Pediatricians

Shift

ABSTRACT

Introduction: People with a reduced nighttime dip in blood pressure have an increased cardiovascular risk. Our objective was to describe the different patterns in blood pressure (BP) among pediatricians who work in long on-duty shifts in relation with sex, medical rank and sleeping time.

Methods: Descriptive, cross-sectional, two-center study. On duty pediatric Resident physicians and pediatric Consultants were recruited between January 2018 and December 2021.

Results: Fifty-one physicians were included in the study (78.4% female, 66.7% Resident physicians). Resident physicians had a higher night/day ratio (0.91 vs 0.85; $p < 0.001$) and a shorter nighttime period (3.87 vs 5.41, $p < 0.001$) than Consultants. Physicians sleeping less than 5 h had a higher night/day ratio (0.91 vs 0.87, $p = 0.014$). Being a Resident showed a ~4.5-fold increased risk of having a non-dipping BP pattern compared to Consultants.

Conclusion: We found a potential link between both being a Resident and, probably, having shorter sleeping time, and the non-dipping BP pattern in physicians during prolonged shifts.

© 2022 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: javigongar8@gmail.com (J. Gonzalez-Garcia).

<https://doi.org/10.1016/j.nefro.2022.05.008>

0211-6995/© 2022 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Patrón no-dipper en la presión arterial de médicos pediatras durante las guardias presenciales

RESUMEN

Palabras clave:

Hipertensión
No dipper
Pediatras
Guardia

Introducción: Las personas con un descenso nocturno reducido de la presión arterial tienen mayor riesgo cardiovascular. Nuestro objetivo fue describir los diferentes patrones de presión arterial en los pediatras que trabajan de guardia con presencia física, en relación con el sexo, la categoría profesional y el tiempo de sueño.

Métodos: Se realizó un estudio descriptivo, transversal, bicéntrico. Se reclutó a médicos residentes y adjuntos de pediatría, de guardia con presencia física, entre enero de 2018 y diciembre de 2021.

Resultados: Fueron incluidos en el estudio 51 médicos (78,4% mujeres; 66,7% médicos residentes). Los médicos residentes presentaron un cociente de presión arterial noche/día mayor (0,91 vs. 0,85; $p < 0,001$) y un tiempo de sueño menor (3,87 vs. 5,41; $p < 0,001$) que los adjuntos. Los participantes que durmieron menos de 5 horas presentaron un cociente de presión arterial noche/día mayor (0,91 vs. 0,87; $p = 0,014$). Ser médico residente demostró tener aumentado el riesgo de presentar un patrón no dipper en más de 4,5 veces respecto a los médicos adjuntos.

Conclusiones: Encontramos un vínculo potencial entre ser médico residente y, probablemente, tener menos horas de sueño, y el patrón de no descenso nocturno de la presión arterial en los médicos durante las guardias de presencia física.

© 2022 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Hypertension (HTN) is a major preventable cause of cardiovascular diseases and all-cause mortality worldwide, with an overall prevalence of 30–45%.¹ There is growing evidence suggesting that shift work is a risk factor for many negative health outcomes, including cardiovascular disease and HTN.^{2–4} Also, there is evidence noting an association between shift work and sleep deprivation, and between sleep deprivation and HTN.^{5–7}

So far, most studies about shift work classified HTN based on office blood pressure (BP) measurement.⁴ However, ambulatory blood pressure monitoring (ABPM) techniques have expanded the knowledge regarding the circadian rhythm of BP and now, ABPM is considered as the gold standard method for evaluating BP, providing a more accurate estimation of BP related risks.^{8,9} People with a reduced nighttime dip in BP (<10% of the daytime average BP or a night/day ratio > 0.9) have an increased cardiovascular risk.^{10,11}

The main objective of the study was to describe the different patterns in BP among pediatricians who work in long on-duty shifts in relation with sex, medical rank (resident physicians vs consultants) and sleeping time.

Methods

Study design

We conducted a descriptive, cross-sectional, two-center study. On duty pediatric Resident physicians and pediatric Consultants working in two hospitals were recruited between January

2018 and December 2021. One center was a tertiary teaching hospital and the other one was a secondary teaching hospital, both in Spain. Physicians with conditions that may affect their fluid and electrolyte balance (impaired kidney function, pregnancy, diabetes, hypertension) and those who refused to participate in the study were excluded. Weight and height were measured for each subject and total body surface (TBS) and body mass index (BMI) were calculated. Sex, age and relevant past medical history was collected for each subject. During the 17-h shift, hours of activity, hours of rest and ABPM were recorded. Each participant documented the hours of active work and the hours of sleep during the shift.

BP measurement was carried out following the recommendations of the European Society of Hypertension.¹⁰ Basal systolic BP (SBP) and diastolic BP (DBP) were measured at the office with a validated automatic oscillometric device (SunTech CT40, SunTech Medical Inc., Morrisville, NC, USA), after 5-min rest in a sitting position. BP values were estimated as the mean of two readings. Thereafter, 17-h ABPM was performed using the SpaceLabs 90207 automated non-invasive oscillometric device (Spacelabs Medical, Space-Labs Inc., Redmond, WA, USA), programmed to register BP every 30-min from 11 pm to 7 am and every 20-min the rest of the time. We selected cuff size to fit the individual's arm circumference according to the device's instructions. Pediatricians were instructed to maintain their usual activities on duty, return the following morning for device removal and keep the arm extended and immobile at the time of each cuff inflation, if it was possible. Valid registries had to fulfill a series of pre-established criteria, including 80% of systolic BPs and diastolic BPs successful recordings during the daytime and nighttime periods, 17-h duration, and at least one BP measure-

Table 1 – Characteristics of population according to sex. Tables shows means \pm SD or median (p25–p75) according to the test used for comparison.

	Male	Female	p value
N	11	40	–
Age (years)	29 (27.5–40.5)	28 (26–30)	0.206
Weight (kg)	71.2 \pm 7.1	58.5 \pm 7.1	<0.001
BMI (kg/m ²)	23.1 \pm 2.4	22.2 \pm 2.2	0.227
TBS (m ²)	1.86 \pm 0.1	1.62 \pm 0.12	<0.001
Nighttime period (h)	5 (5–5)	4 (3–5)	0.076
Basal SBP (mmHg)	125.2 \pm 6.6	117.9 \pm 12	0.013
Basal DBP (mmHg)	76.3 \pm 7.1	74.4 \pm 7.9	0.469
Daytime SBP (mmHg)	125.4 \pm 8.4	114.2 \pm 8.7	<0.001
Daytime DBP (mmHg)	77.9 \pm 6.9	75.5 \pm 5.8	0.249
Nighttime SBP (mmHg)	109.2 \pm 12.1	103.1 \pm 8.6	0.063
Nighttime DBP (mmHg)	60.6 \pm 7.3	62 \pm 6.6	0.554
24 h HR (bpm)	66.9 \pm 13.8	75.7 \pm 8.5	0.066
Daytime HR (bpm)	68.9 \pm 14.3	77.8 \pm 8.5	0.073
Nighttime HR (bpm)	56 \pm 11.4	65.1 \pm 10.4	0.016
Night/day ratio SBP (mmHg)	0.87 \pm 0.07	0.9 \pm 0.05	0.095
Night/day ratio DBP (mmHg)	0.77 \pm 0.07	0.82 \pm 0.07	0.044

BMI: body mass index; bpm: beats per minute; DBP: diastolic blood pressure; HR: heart rate; mmHg: millimeters of mercury; SBP: systolic blood pressure; TBS: total body surface.

ment per hour. Daytime and nighttime periods were defined individually according to the hours of active work and sleep, respectively.

Definitions

- Night/day ratio: ratio of average nighttime BP to average daytime BP for SBP and DBP separately.
- Definition of Circadian Patterns: $\geq 10\%$ reduction in the average SBP during the nighttime period compared to daytime mean SBP was considered normal dipping pattern (dipper). Contrariwise, a $< 10\%$ reduction of mean nighttime SBP compared to daytime mean SBP was considered abnormal dipping pattern (non-dipper).

Statistical analysis

Statistical analysis was carried out using the R program (R Development Core Team), version 3.6.3. Descriptive analysis was performed providing relative and absolute frequency distributions for qualitative variables, and measures of position and dispersion for the quantitatives. Normally distributed data were expressed as mean \pm SD, non-normally distributed data as median and interquartile range and categorical variables were reported as percentages. Differences according to sex, medical rank and hours of sleep, grouped into two categories, were assessed with the Student's t test or Wilcoxon for independent samples, after verification of the normality hypothesis. Univariate and multivariate logistic regression models were built to determine the dipping BP pattern profile. A p value less than 0.05 was considered statistically significant.

We obtained approval from the Clinical Research Ethics Committee of the Autonomous Region of Asturias. All persons gave their written informed consent prior to their inclusion in the study.

Results

Fifty-one physicians were included in the study (mean age 31 ± 7.5 years old). Forty (78.4%) were female, 34 were resident physicians (66.7%). Mean of BMI was 22.3 ± 2.2 . All ABPM records were valid. Description of the population according to sex appears in [Table 1](#).

Resident physicians had a higher night/day ratio (0.91 ± 0.05 vs 0.85 ± 0.05 ; $p < 0.001$) and a shorter nighttime period (3.87 ± 1.36 h vs 5.41 ± 1.23 h, $p < 0.001$) than Consultants. Description of the population according to medical rank appears in [Table 2](#), while [Table 3](#) describes the population according to hours of sleep. Physicians who sleep less than 5 h had a higher night/day ratio (0.91 ± 0.04 vs 0.87 ± 0.07 , $p = 0.014$).

Univariate and multivariate logistic regression models were built to determine the dipper profile ([Table 4](#)). We found a significant relationship between medical rank and dipping BP pattern, which remains after adjusting for sleeping time. Being a Resident seems to increase the probability of non-dipping BP pattern more than 4.5 times compared to being a Consultant ($p = 0.033$).

Discussion

The present study describes a potential link between being a Resident and the non-dipping BP pattern in physicians during prolonged shifts.

So far, recognized reasons for an absence of nocturnal BP dipping are sleep disturbance, obstructive sleep, apnea, obesity, high salt intake in salt-sensitive subjects, orthostatic hypotension, autonomic dysfunction, chronic kidney disease, diabetic neuropathy and old age.¹² However, none of the participants of our study met these characteristics, except sleep disturbance during the shift.

Table 2 – Characteristics of population according to medical rank. Tables shows means \pm SD or median (p25–p75) according to the test used for comparison.

	Residents	Consultants	p value
N	34	17	–
Age (years)	27 (26–28)	37 (32–41)	<0.001
Weight (kg)	60.1 \pm 9.2	63.6 \pm 7.7	0.314
BMI (kg/m ²)	22.4 \pm 2.4	22.5 \pm 1.9	0.951
TBS (m ²)	1.65 \pm 0.16	1.72 \pm 0.13	0.269
Nighttime period (h)	4 (3–5)	5 (5–6)	<0.001
Basal SBP (mmHg)	118.6 \pm 12.3	121.4 \pm 9.3	0.409
Basal DBP (mmHg)	75.5 \pm 8.3	73.3 \pm 6.3	0.339
Daytime SBP (mmHg)	115.5 \pm 10.5	118.8 \pm 7.8	0.254
Daytime DBP (mmHg)	76.1 \pm 7	75.9 \pm 3.6	0.875
Nighttime SBP (mmHg)	105.6 \pm 9.7	101.9 \pm 9.5	0.204
Nighttime DBP (mmHg)	62.7 \pm 7.5	59.7 \pm 4.3	0.075
24 h HR (bpm)	76.2 \pm 8.7	69.1 \pm 11.9	0.02
Daytime HR (bpm)	78.1 \pm 8.9	71.3 \pm 2.3	0.029
Nighttime HR (bpm)	64.6 \pm 10.7	60.1 \pm 11.7	0.18
Night/day ratio SBP (mmHg)	0.91 \pm 0.05	0.85 \pm 0.05	<0.001
Night/day ratio DBP (mmHg)	0.83 \pm 0.07	0.78 \pm 0.05	0.031

BMI: body mass index; bpm: beats per minute; DBP: diastolic blood pressure; HR: heart rate; mmHg: millimeters of mercury; SBP: systolic blood pressure; TBS: total body surface.

Table 3 – Characteristics of population according to hours of sleep. Tables shows means \pm SD or median (p25–p75) according to the test used for comparison.

	<5 h of sleep	>5 h of sleep	p value
N	23	28	–
Residents (N)	21	13	–
Consultants (N)	2	15	–
Age (years)	28 (26–29)	29.5 (27–39)	0.003
Weight (kg)	59.3 \pm 7.5	62.8 \pm 9.6	0.165
BMI (kg/m ²)	22.4 \pm 1.7	22.3 \pm 2.5	0.887
TBS (m ²)	1.64 \pm 0.15	1.70 \pm 0.16	0.196
Basal SBP (mmHg)	120.2 \pm 13	118.9 \pm 10	0.679
Basal DBP (mmHg)	76.5 \pm 8.8	73.4 \pm 6.5	0.147
Daytime SBP (mmHg)	116.1 \pm 10.4	117 \pm 9.3	0.733
Daytime DBP (mmHg)	76.4 \pm 7.3	75.7 \pm 4.8	0.721
Nighttime SBP (mmHg)	106.4 \pm 10.1	102.7 \pm 9.1	0.184
Nighttime DBP (mmHg)	63.7 \pm 7.8	60 \pm 5.2	0.059
24 h HR (bpm)	77.9 \pm 8.2	70.5 \pm 10.9	0.009
Daytime HR (bpm)	79.5 \pm 8.6	72.9 \pm 11.1	0.024
Nighttime HR (bpm)	68.1 \pm 8.9	59 \pm 11.3	0.003
Night/day ratio SBP (mmHg)	0.91 \pm 0.04	0.87 \pm 0.07	0.014
Night/day ratio DBP (mmHg)	0.84 \pm 0.06	0.79 \pm 0.07	0.011

BMI: body mass index; bpm: beats per minute; DBP: diastolic blood pressure; HR: heart rate; mmHg: millimeters of mercury; SBP: systolic blood pressure; TBS: total body surface.

Table 4 – Univariate and multivariate analysis to determine dipping BP pattern profile.

	Dipping pattern		Univariate OR	Multivariate OR
	Non-dipper	Dipper		
Medical rank				
Residents	22 (64.7%)	12 (35.3%)	–	–
Consultants	4 (23.5%)	13 (76.5%)	5.96 (1.70–25.07, p = 0.008)	4.86 (1.20–23.13, p = 0.033)
Hours of sleep				
<5 h	15 (65.2%)	8 (34.8%)	–	–
>5 h	11 (39.3%)	17 (60.7%)	2.90 (0.94–9.47, p = 0.069)	1.54 (0.40–5.73, p = 0.522)

OR: odds ratio.

Furthermore, non-dipping pattern was more frequent in Resident physicians (who were under 30 years of age), in contrast to previous studies showing an increased risk of a non-dipping BP related to an increasing age.¹² This could be related to the fact that the study was carried out in university hospitals, where Resident physicians have increased workload because they are responsible for providing initial care to the patients.

It has been reported a significantly higher prevalence of HTN in people with sleep disturbances, such as airline pilots.¹³ Sleep disturbance has been described as the main cause of nocturnal hypertension.¹⁴ Anxiety disorder¹⁵ and insomnia¹⁶ have also been associated to nocturnal hypertension, probably resulted from excessive tension due to sleep disturbance during the night. However, we have not found a significant relationship between non-dipping BP pattern in Resident physicians, based on sleeping time, after adjusting in the multivariate analysis. Future studies with larger number of subjects are needed to determine if sleeping time influences the BP pattern in Resident physicians.

It has also been reported an association between history of shift work and rates of hypertension in both men and women.⁷ We have found that females were classified as a non-dipping pattern more frequently than males, but they were no differences by sex in night/day ratio SBP. Also, in our study, males and females had normal BP values, based on classification of hypertension grade¹ and differences by sex are similar to those of the general population.

Some studies^{17,18} reported that circumstances which cause frequent awakenings during sleep increased heart rate (HR). Although we found that heart rate values are higher in individuals who sleep fewer hours, resting values do not exceed 80 beats/min, so we cannot consider it an added cardiovascular risk factor.¹

Our study has several limitations. First, it is a cross-sectional study and results are limited to a low number of subjects of Caucasian population in a single Health area in the Autonomous Region of Asturias, in Northern Spain. Second, results are limited to a single time on each participant. Dipping status has a low reproducibility, with up to 40% of individuals from Europe¹⁹ changing status between repeated recordings.

In contrast, this is one of the few times that continuous BP measurement and non-dipping pattern are reported during shift work and the first time that BP is reported during a medical shift in a pediatric department, which could have implications for future studies on the impact of medical shifts on the health of the physicians. Moreover, the main strength of this study is that it was performed in a well-delimited area, which allows a detailed study of each participant.

Conclusions

We found a potential link between both being a Resident and, probably, having shorter sleeping time, and the non-dipping blood pressure pattern in physicians during prolonged shifts. Future studies with larger numbers of subjects are needed to corroborate the association between non-dipping pattern and shift work in medical shifts and obtain stronger conclusions.

Conflict of interest

The authors have no conflicts of interest to disclose.

Funding source

This research has not received specific aid from public sector agencies, the commercial sector or non-profit entities.

Financial disclosure

The authors have no financial relationships relevant to this article to disclose.

Ethics

We obtained approval from the Clinical Research Ethics Committee of the Autonomous Region of Asturias. All persons gave their written informed consent prior to their inclusion in the study.

All authors have read and approved the final revised version. They are willing to take responsibility for the entire manuscript. All authors substantially contributed to the writing (i.e. drafting and/or critical revision) of the manuscript.

Acknowledgments

We would like to thank Statistical Consulting Unit of the Scientific-Technical Services of the University of Oviedo for their cooperation in the statistical analysis.

REFERENCES

1. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, et al. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J*. 2018;39:3021-104.
2. Rivera AS, Akanbi M, O'Dwyer LC, McHugh M. Shift work and long work hours and their association with chronic health conditions: a systematic review of systematic reviews with meta-analyses. *PLoS One*. 2020;15:e0231037.
3. Puttonen S, Härmä M, Hublin C. Shift work and cardiovascular disease – pathways from circadian stress to morbidity. *Scand J Work Environ Health*. 2010;36:96-108.
4. Manohar S, Thongprayoon C, Cheungpasitporn W, Mao MA, Herrmann SM. Associations of rotational shift work and night shift status with hypertension: a systematic review and meta-analysis. *J Hypertens*. 2017;35:1929-37.
5. Gottlieb DJ, Redline S, Nieto FJ, Baldwin CM, Newman AB, Resnick HE, et al. Association of usual sleep duration with hypertension: the Sleep Heart Health Study. *Sleep*. 2006;29:1009-14.
6. Akerstedt T. Shift work and disturbed sleep/wakefulness. *Sleep Med Rev*. 1998;2:117-28.
7. Rahim A, McIsaac MA, Aronson KJ, Smith PM, Tranmer JE. The associations of shift work, sleep quality, and incidence of hypertension in ontario adults: a population-based study. *Can J Cardiol*. 2021;37:513-8.
8. Gorostidi M, Sobrino J, Segura J, Sierra C, de la Sierra A, Hernández del Rey R, et al. Ambulatory blood pressure

- monitoring in hypertensive patients with high cardiovascular risk: a cross-sectional analysis of a 20,000-patient database in Spain. *J Hypertens.* 2007;25:977–84.
9. Metoki H, Ohkubo T, Imai Y. Diurnal blood pressure variation and cardiovascular prognosis in a community-based study of Ohasama, Japan. *Hypertens Res.* 2010;33:652–6.
 10. Stergiou GS, Palatini P, Parati G, O'Brien E, Januszewicz A, Lurbe E, et al. European Society of Hypertension Council and the European Society of Hypertension Working Group on Blood Pressure Monitoring and Cardiovascular Variability. 2021 European Society of Hypertension practice guidelines for office and out-of-office blood pressure measurement. *J Hypertens.* 2021;39:1293–302.
 11. Boggia J, Li Y, Thijs L, Hansen TW, Kikuya M, Björklund-Bodegård K, et al. International Database on Ambulatory blood pressure monitoring in relation to Cardiovascular Outcomes (IDACO) investigators. Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. *Lancet.* 2007;370:1219–29.
 12. de la Sierra A, Redon J, Banegas JR, Segura J, Parati G, Gorostidi M, et al. Prevalence and factors associated with circadian blood pressure patterns in hypertensive patients. *Hypertension.* 2009;53:466–72.
 13. Wilson D, Driller M, Johnston B, Gill N. The prevalence and distribution of health risk factors in airline pilots: a cross-sectional comparison with the general population. *Aust N Z J Public Health.* 2022.
 14. Henskens LH, van Boxtel MP, Kroon AA, van Oostenbrugge RJ, Lodder J, de Leeuw PW. Subjective sleep disturbance increases the nocturnal blood pressure level and attenuates the correlation with target-organ damage. *J Hypertens.* 2011;29:242–50.
 15. Kayano H, Koba S, Matsui T, Fukuoka H, Toshida T, Sakai T, et al. Anxiety disorder is associated with nocturnal and early morning hypertension with or without morning surge: ambulatory blood pressure monitoring. *Circ J.* 2012;76:1670–7.
 16. Ostroumova TM, Parfenov VA, Ostroumova OD, Kochetkov AI. Hypertension and insomnia. *Ter Arkh.* 2020 Jan 15;92:69–75.
 17. Boudreau P, Yeh WH, Dumont GA, Boivin DB. Circadian variation of heart rate variability across sleep stages. *Sleep.* 2013;36:1919–28.
 18. Sforza E, Pichot V, Cervena K, Barthélémy JC, Roche F. Cardiac variability and heart-rate increment as a marker of sleep fragmentation in patients with a sleep disorder: a preliminary study. *Sleep.* 2007;30:43–51.
 19. Omboni S, Parati G, Palatini P, Vanasia A, Muiesan ML, Cuspidi C, et al. Reproducibility and clinical value of nocturnal hypotension: prospective evidence from the SAMPLE study. *J Hypertens.* 1998;16:733–8.