



## Original article

# Clinicopathological characteristics and outcomes of patients with unexplained renal impairment: A single center study

Q1 Radwa A. Ellisy <sup>a,\*</sup>, Effat A. Tony <sup>a</sup>, Wesam Ismail <sup>b</sup>, Rabab Radi <sup>c</sup>,  
Mohamed Ismail Seddik <sup>d</sup>, Essam M. Abdel Aziz <sup>a</sup>

<sup>a</sup> Division of Nephrology, Internal Medicine Department, Faculty of Medicine, Assiut University, Assiut, Egypt

<sup>b</sup> Department of Pathology, Faculty of Medicine, Beni-Suef University, Beni-Suef, Egypt

<sup>c</sup> Department of Pathology, Faculty of Medicine, Cairo University, Giza, Egypt

<sup>d</sup> Clinical Pathology Department, Faculty of Medicine, Assiut University, Assiut, Egypt

## ARTICLE INFO

## Article history:

Received 25 May 2024

Accepted 20 October 2024

Available online xxx

## Keywords:

Renal biopsy

Unexplained renal impairment

CKDu

Histopathology

## ABSTRACT

**Background:** Chronic kidney disease (CKD) is a growing health problem. About 20% of CKD patients have undetermined causes. Data describing histopathological patterns of unexplained impaired kidney functions in Egypt are lacking. We aimed to identify the clinicopathological characteristics and short-term outcomes of adult cases with unexplained impaired kidney functions.

**Methods:** We conducted a prospective study in Assiut University Hospital from August 2018 to May 2022. It included patients with unexplained elevated serum creatinine (serum creatinine > 115  $\mu\text{mol/L}$ ) who underwent renal biopsies. Descriptive statistics were used to summarize data on patient demographics, histopathological patterns, and outcomes after three months. Clinical trial registration number: NCT03586531.

**Results:** Overall, 210 native renal biopsies were included in the analysis. Glomerular diseases were the most common pathological finding ( $n = 88$ , 44.9%), amyloidosis and FSGS were the most prevalent glomerular pathology (15.2%, 14.3%, respectively). Chronic kidney disease of unknown etiology (CKDu) was diagnosed in 8.1% of the cases; histology suggestive of genetic origin was found in 2.5%, and LECT amyloidosis was found in 3.8% of the cases. Poor outcomes were observed in 43.6% of the patients i.e., renal replacement therapy or death. Treatment strategies were changed based on the biopsy findings in 86 patients (40%).

**Conclusion:** Amyloidosis and FSGS were the most common causes of unexplained renal impairment. CKDu is not uncommon in Egypt, and more preventive measures are needed. This study supports the irreplaceable role of renal biopsy in disease diagnosis, treatment decisions, and predicting prognosis even in advanced stages.

© 2024 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [radwa.ellisy15@aun.edu.eg](mailto:radwa.ellisy15@aun.edu.eg) (R.A. Ellisy).

<https://doi.org/10.1016/j.nefro.2024.10.006>

0211-6995/© 2024 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Ellisy R,A, et al. Clinicopathological characteristics and outcomes of patients with unexplained renal impairment: A single center study. Nefrologia. 2024. <https://doi.org/10.1016/j.nefro.2024.10.006>

## Características clinicopatológicas y resultados de pacientes con insuficiencia renal inexplicable: un estudio de un solo centro

### R E S U M E N

#### Palabras clave:

Biopsia renal  
Insuficiencia renal inexplicable  
ERCn  
Histopatología

**Antecedentes:** La enfermedad renal crónica (ERC) es un problema de salud en crecimiento. Aproximadamente el 20% de los pacientes con ERC tienen causas no determinadas. Los datos que describen los patrones histopatológicos de la disfunción renal inexplicable en Egipto son escasos. Nuestro objetivo fue identificar las características clinicopatológicas y los resultados a corto plazo de casos adultos con disfunción renal inexplicable.

**Métodos:** Realizamos un estudio prospectivo en el Hospital Universitario de Assiut desde agosto de 2018 hasta mayo de 2022. Se incluyeron pacientes con niveles elevados inexplicados de creatinina sérica (creatinina sérica > 115  $\mu\text{mol/l}$ ) que se sometieron a biopsias renales. Se utilizaron estadísticas descriptivas para resumir los datos sobre la demografía de los pacientes, los patrones histopatológicos y los resultados después de tres meses. Número de registro del ensayo clínico: NCT03586531.

**Resultados:** En total, se incluyeron 210 biopsias renales nativas en el análisis. Las enfermedades glomerulares fueron el hallazgo patológico más común ( $n = 88$ , 44,9%), siendo la amiloidosis y la glomeruloesclerosis focal y segmentaria (FSGS) las patologías glomerulares más prevalentes (15,2% y 14,3%, respectivamente). La enfermedad renal crónica de causa desconocida (ERCn) se diagnosticó en el 8,1% de los casos; la histología sugerente de origen genético se encontró en el 2,5%, y la amiloidosis LECT se detectó en el 3,8% de los casos. Se observaron malos resultados en el 43,6% de los pacientes, es decir, terapia de reemplazo renal o muerte. Las estrategias de tratamiento se modificaron en 86 pacientes (40%) basándose en los hallazgos de la biopsia.

**Conclusión:** La amiloidosis y la FSGS fueron las causas más comunes de insuficiencia renal inexplicable. La ERCn no es infrecuente en Egipto, y se necesitan más medidas preventivas. Este estudio respalda el papel insustituible de la biopsia renal en el diagnóstico de enfermedades, las decisiones de tratamiento y la predicción del pronóstico, incluso en etapas avanzadas.

© 2024 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licencias/by-nc-nd/4.0/>).

## Introduction

Chronic kidney disease (CKD) is a growing healthcare problem that affects more than 800 million people worldwide. Its rank is projected to advance from the 16th to the 5th cause of death globally by 2040.<sup>1</sup> CKD is mostly caused by diabetes mellitus (DM), hypertension (HTN), and glomerular diseases. However, sometimes the etiology of CKD is unknown owing to the absence of identifiable risk factors, or specific histological findings.<sup>2</sup> This condition is called chronic kidney disease of unknown etiology (CKDu).<sup>3</sup>

In a 2006 study, CKDu was reported in 27% of patients who received dialysis in El-Minia Governorate of Egypt; however, their diagnosis was not based on histological findings.<sup>4</sup> It was uncertain if the patients developed their condition after exposure to risk factors related to the agricultural community or if they presented late after having undiagnosed treatable conditions. A definite diagnosis can be achieved through renal biopsy, especially when conventional workup is inconclusive. Renal biopsies also provide prognostic information for renal and patient survival and can help in deciding treatment options and pre-transplant workup, even in advanced cases.<sup>5</sup>

In an Egyptian study, researchers analyzed the clinic pathological data of patients who underwent renal biopsy at their center in 2020. They found that 47.8% of the biopsies were performed for patients with accidentally discovered renal impairment.<sup>6</sup> However, to date, there is no Egyptian registry for data on the pathological results of renal biopsies in patients with unexplained renal impairment. Identifying the underlying pathology in these patients could help healthcare professionals provide optimum and timely therapeutic options and preventive measures for their condition. Herein, we aimed to identify the histopathological pattern of patients with unexplained renal impairment, their clinical outcomes, and the clinic pathological characteristics of patients with CKDu.

## Materials and methods

### Study design

This prospective observational study was conducted at the Nephrology Unit of Assiut University Hospital, which provides tertiary care for patients from Upper Egypt. It included adult

90 patients with renal impairment that could not be diagnosed  
91 with clinical and laboratory parameters (unexplained renal  
92 impairment). Patients were recruited between August 2018  
93 and May 2022 and followed up for three months after the  
94 date of their biopsy. The study protocol was approved by the  
95 Faculty of Medicine Ethical Review Board at Assiut University  
96 (Approval number: IRB17200010), and registered in clinical-  
97 trials.gov, registration number: NCT03586531. The study was  
98 performed in accordance with the principles of the Declara-  
99 tion of Helsinki.

## 100 Patients

101 Informed consent was obtained from all individual par-  
102 ticipants included in the study. All patients fulfilled the  
103 following inclusion criteria: age  $\geq$  18 years old, elevated serum  
104 creatinine  $>$  115  $\mu$ mol/L, negative serological markers (ANA,  
105 antidsDNA, perinuclear or cytoplasmic antineutrophil cyto-  
106 plasmic antibodies, and normal complement (C3 and C4))  
107 levels. Patients with the following criteria were excluded from  
108 the study: diabetes with clinical and laboratory data favoring  
109 diabetic nephropathy,<sup>7</sup> hypertension with clinical data sug-  
110 gestive of hypertensive nephrosclerosis,<sup>8</sup> suspicion of renal  
111 involvement in systemic disease, history of nephrotoxic drugs,  
112 the presence of relative or absolute contraindications for the  
113 procedure, kidney size  $<$  8 cm, and pregnancy.

## 114 Methods

115 Patients' data were collected at the biopsy visit and included  
116 demographics, detailed medical and medication history, and  
117 essential laboratory investigations. Serum creatinine was  
118 measured using a Jaffe-based method, and urine proteins  
119 were measured by an immunoturbidometric method. GFR was  
120 estimated using the creatinine-based Chronic Kidney Disease  
121 Epidemiology Collaboration equation.

## 122 Biopsy procedure

123 Biopsies were done by skilled nephrologists according to  
124 standard procedures.<sup>9</sup> Biopsy complications were catego-  
125 rized as minor and major complications. Minor complications  
126 included flank pain, hematuria, and/or spontaneously resolv-  
127 ing hematoma without the need for further intervention.  
128 Major complications were those that resulted in hemody-  
129 namic instability, need for an intervention, such as transfusion  
130 of blood products, further imaging, invasive radiologic or sur-  
131 gical procedures, or death.

132 All biopsies were processed routinely for light microscopy  
133 and immunohistochemistry and examined by one of two  
134 experts in nephropathology. Immunoperoxidase (IP) studies  
135 were performed using antihuman IgG, IgA, IgM, C3, kappa,  
136 and lambda light chains. Cases were examined by electron  
137 microscopy (EM) when feasible based on the findings of clinical  
138 examination.

## 139 Treatment

140 Treatment plans were either "specific treatment," which were  
141 dependent or modified according to the histological diagnosis,

or "supportive treatment" due to the presence of advanced  
142 lesions or the unavailability of specific treatment. 143

## 144 Patients' outcomes

145 Patients were followed for 3 months for the following end  
146 points: renal recovery (creatinine return to/or within 25% of  
147 the baseline value and dialysis independence),<sup>10</sup> need for  
148 maintenance dialysis, or death from causes not related to the  
149 biopsy procedure.

## 150 Statistics

151 Data analysis was performed using SPSS version 26.0 (SPSS  
152 Inc., Chicago, IL, US). Means  $\pm$  SD were used to describe nor-  
153 mally distributed variables, while medians and ranges were  
154 used to describe non-normally distributed variables. Chi-  
155 square/Fisher Exact tests were used to compare proportions  
156 between groups. One way ANOVA or Kruskal-Wallis tests  
157 were used to compare mean and median differences between  
158 groups. A post hoc test for pairwise comparison with Bon-  
159 ferroni correction was done to identify significance between  
160 more than two groups. Univariate logistic regression analysis  
161 was performed to identify possible predictors for poor out-  
162 comes among patients, and significant variables were entered  
163 in a multivariate backward Wald logistic regression analysis  
164 to calculate adjusted odds ratios (AOR). Statistical significance  
165 was set at  $p <$  0.05.

## 166 Results

167 During the study period, 210 of 687 native biopsies (30.6%)  
168 were performed for unexplained renal impairment. Two were  
169 excluded from the analysis due to inadequate sampling  
170 (medulla only). The patients were mostly middle-aged with  
171 near equal sex distribution. Their mean serum creatinine  
172 level was seven folds higher than normal. The most frequent  
173 comorbidities were hypertension followed by diabetes. Post-  
174 biopsy complications occurred in 19 patients. Six patients  
175 developed major complications in the form of large per-  
176 inepric hematoma necessitating blood transfusion. Five of  
177 these patients were managed conservatively, and one under-  
178 went super-selective renal artery angioembolization to control  
179 bleeding. The other 13 patients had minor complications as a  
180 small perinepric hematoma (five patients) and gross hema-  
181 turia (eight patients) (Table 1).

## 181 Histological diagnosis

182 Patients were categorized according to their histological diag-  
183 nosis into five groups: glomerular, tubulointerstitial, vascular,  
184 paraprotein-related renal disease (PRDs), and pathology sug-  
185 gestive of CKDu (Fig. 1). The most frequent diagnoses were  
186 glomerular and tubulointerstitial diseases, accounting for  
187 two-thirds of all biopsies. The most frequent glomerular  
188 pathologies were amyloidosis ( $n=32$ , 15.2%) and focal seg-  
189 mental glomerulosclerosis (FSGS, 14.3%). In the amyloidosis  
190 group, 14 patients had ALECT2; 18 patients had AA amy-  
191 loidosis; and two patients had AL amyloidosis (categorized


**Table 1 – Demographic and clinical data of the studied participants (n = 210).**

Demographic data	
Age, year	43.7 ± 14.9
<40	88 (41.9%)
40 to <65	95 (45.2%)
≥65	27 (12.9%)
Male	
Current smoking	51 (24.3%)
Substance abuse	23 (11%)
Farmer	51 (24.3%)
Medical history	
Hypertension	64 (30.5%)
DM	19 (9%)
Cardiac diseases	6 (2.9%)
Chronic kidney disease	5 (2.4%)
Thyroid diseases	4 (1.9%)
COPD	1 (0.5%)
Clinical data	
SBP, mmHg	134 ± 14.9
DBP, mmHg	83.2 ± 8.3
MAP, mmHg	100.1 ± 9.8
Nephrotic syndrome	46 (21.9%)
Creatinine, μmol/l	724 (136–2300)
Microscopic hematuria	88 (41.9%)
24-h protein, mg/day	1217 (117–9700)
eGFR (ml/min/m <sup>2</sup> ) <sup>a</sup>	6 (2–50)
Lt kidney length, cm	10.1 ± 1.4

Data are expressed as percentages, (mean ± SD), or (median, range) as appropriate.

COPD: Chronic Obstructive Pulmonary Disease, eGFR: estimated glomerular filtration rate, DBP: diastolic blood pressure, DM: diabetes mellitus, MAP: mean arterial blood pressure, SBP: systolic blood pressure.

<sup>a</sup> eGFR was estimated by CKD-EPI equation.



**Fig. 1 – histological groups and subgroups of the studied groups. AG: advanced glomerulosclerosis, AIN: acute interstitial nephritis, AL: light chain amyloidosis, ATN: acute tubular necrosis, Cast N: cast nephropathy, CKDu: chronic kidney disease of undetermined etiology, CNS TIN: chronic non-specific tubulointerstitial nephritis, CTIN: chronic tubulointerstitial nephritis, FSGS: focal and segmental glomerulosclerosis, GDs: glomerular diseases, MIDDs: monoclonal immunoglobulin deposition disease; MN: membranous nephropathy, MPGN: membranoproliferative pattern glomerulonephritis, NGN: necrotizing glomerulonephritis, PRDs: paraprotein-related renal diseases, TIDs: tubulointerstitial diseases, TMA: thrombotic microangiopathy, VDs: vascular diseases.**

as PRDs). Within the tubulointerstitial disease (TIDs) group, five patients (2.4%) had histological criteria highly suggestive of genetically inherited diseases: autosomal dominant tubulointerstitial kidney disease (ADTKD) in three patients, and nephronophthisis in two patients. Eight patients had chronic tubulointerstitial nephritis (3.8%), of whom six had granulomatous nephritis, and two had IgG4 tubulointerstitial nephritis. PRD comprised 12.9% of cases, with light chain cast nephropathy as the predominant type, comprising 10.5% of all cases. Some histological diagnoses are shown in Fig. 2.

### Clinical variables and histopathological characteristics

We compared the five main histological groups as regards their clinical variables and compared each histological group with the CKDu group. The youngest patients were in the CKDu group, and the oldest patients were in the PRDs group, with a mean age of  $37.8 \pm 13.6$  vs.  $50.5 \pm 9.7$  years ( $p = 0.006$ ), respectively. The proportion of males was highest in the CKDu group (11/17, 64.7%) and lowest in the TIDs group (11/17, 64.7% vs. 15/43, 34.9%;  $p = 0.036$ , respectively). The greatest proportion of patients working as farmers were in the CKDu group and comprised 47.1% of patients ( $n = 8$ ,  $p = 0.001$ ) (Table 2).

Forty-six patients (21.9%) presented with renal impairment and nephrotic syndrome, unsurprisingly most of them had glomerular diseases and none of them had TIDs (Table 2).


The lowest mean serum creatinine level was observed in the glomerular disease group ( $638.5 \mu\text{mol/L}$ ), while the highest level was observed in the tubulointerstitial diseases group ( $975 \mu\text{mol/L}$ ), and the difference was statistically significant (Table 2).

The median iPTH was high in all groups, with a statistically significant difference between the CKDu group and other groups. Sixty percent of the patients presented with sub-nephrotic range proteinuria (Fig. 3).

Advanced chronic changes, such as glomerular sclerosis, tubular atrophy and interstitial fibrosis were more frequently noted in the CKDu group compared with other groups ( $p = 0.033$ ,  $p = 0.002$ , and  $p < 0.001$ , respectively) (Fig. 4).

### Treatment strategies and outcomes

Patients with PRDs or VDs were more likely to be assigned specific treatment based on their biopsy results. However, all



**Fig. 2** – some histological diagnoses. (a) S: under light microscopy using H&E stain showing a segmental area of collapsed capillaries with increased mesangial matrix (original magnification  $\times 200$ ). (b) Picture suggestive of CKDu showing chronic interstitial nephritis, marked fibrosis, and sclerosed glomeruli by PAS stain (original magnification  $\times 200$ ). (c) Picture suggestive of CKDu showing diffuse atrophic changes with mild lymphocytic infiltrate on background of diffuse interstitial fibrosis by Masson trichrome stain (original magnification  $\times 100$ ). (d) Cast nephropathy by H&E, tubules showed moderate injury with large fractured PAS negative hyaline cast surrounded by tubular epithelial cells, inflammatory cells and few giant cells (original magnification  $\times 200$ ). (e) LECT amyloidosis glomerular and interstitial congo red uptake (original magnification  $\times 200$ ). (f) Anti-LECT2 antibody positive reactivity of glomeruli and interstitial deposit (immunoperoxidase technique original magnification  $\times 200$ ).

232 patients in the CKDu group needed only supportive treatment,  
233 including renal replacement therapy (Fig. 5).

234 The patients were followed for 3 months for a more consis-  
235 tent definition for renal disease and proper CKD stages. The  
236 most frequent outcome among CKDu and glomerular disease  
237 groups was the need for maintenance dialysis. Most of the

238 patients who recovered were in the TIDs group. Three patients  
239 died within the follow-up time frame from causes unrelated  
240 to the biopsy procedure (Table 3).

241 Univariate and multivariate regression analysis were per-  
242 formed to determine which factors were associated with  
243 poor outcomes (death and need for maintenance dialysis). A

**Table 2 – Demographic and clinical characteristics of patients who underwent renal biopsy by histological diagnosis.**

Variables	CKDu, n = 17	GDS, n = 88	TIDs, n = 43	PRDs, n = 27	VDs, n = 35	p-Value*
Age, ys	37.8 ± 13.6	44.7 ± 14.6	40.2 ± 16.	50.5 ± 9.7**	43.1 ± 16.3	<b>0.024</b>
Male	11 (64.7%)	53 (60.2%)	15 (34.9%)**	12 (44.4%)	16 (45.7%)	<b>0.048</b>
Farmer	8 (47.1%)	27 (30.7%)	1 (2.3%)**	7 (25.9%)	8 (22.9%)	<b>0.001</b>
Current hypertension	13 (76.5%)	61 (69.3%)	16 (59.3%)	23 (53.5%)	21 (60.0%)	0.310
Nephrotic syndrome	1 (5.7%)	33 (37.5%)**	0 (0%)	5 (18.5%)	7 (20%)	<b>&lt;0.001</b>
SBP, mmHg	140.9 ± 10.4	135.9 ± 16.1	130.7 ± 14.1**	134 ± 14.2	141.3 ± 16.4	0.070
Creatinine (μmol/l)	725 (262–2082)	638.5 (145–2300)	975 (211–1767)	765 (136–1641)	788 (197–2000)	<b>0.044</b>
eGFR ml/min/m <sup>2a</sup>	7 (2–28)	7 (2–50)	7 (2–36)	5 (2–23)	5 (2–39)	0.107
iPTH, pg/ml	300 (50–1900)	214 (9–1652)**	168 (30–1621)**	156 (11–949)**	189 (58–798)**	0.236
Albumin, g/L	29.8 ± 7.1	31.3 ± 6.7	31.9 ± 6.6	28.1 ± 5.8	26.5 ± 8.8	0.058
ALP, U/L	138.6 (78–304)	81.5 (36–176)	79 (29–370)	89 (46–389)	115 (57–214)	0.108
TP, g/L	60.6 ± 7.6	58.4 ± 9.5	64.6 ± 9.9	59.2 ± 9.4	66.6 ± 12.8	0.024
U. ptn, g/24 h	0.8 (0.27–4.33)	2.31 (0.12–9.7)**	0.53 (0.12–3.47)**	1.01 (0.15–4.7)	1.52 (0.15–5.22)	<b>&lt;0.001</b>
Microscopic hematuria	3 (17.6%)	35 (39.8%)	15 (34.9%)	9 (33.3%)	26 (74.3%)**	<b>&lt;0.001</b>
Lt kidney, cm	9.48 ± 0.79	9.88 ± 1.10	10.03 ± 1.21	10.50 ± 1.52**	10.09 ± 1.25	0.065


ALP: alkaline phosphatase, CKDu: chronic kidney disease of unknown etiology, eGFR: estimated glomerular filtration rate, GDS: glomerular diseases, Hgb: hemoglobin level, iPTH: intact parathyroid hormone, PRDs: paraprotein-related renal diseases, TIDs: tubulointerstitial diseases, TP: total protein, VDs: vascular diseases.

<sup>a</sup> eGFR was estimated by CKD-EPI equation.

Data were expressed as frequency (percentage), Mean ± SD, or median (range).

\*  $p < 0.05$  is considered significant. Statistical tests of significance were: one way ANOVA, Kruskal–Wallis, and Chi-square test/Fisher Exact test according to type of data and shown in bold.

\*\* Significant  $p$  after post hoc test for pairwise comparison with Bonferroni correction in comparison to CKDu group.



**Fig. 3 – Proteinuria levels of the studied groups. CKDu: chronic kidney disease of unknown etiology, PRDs: paraprotein-related renal diseases, TIDs: tubulointerstitial diseases, VDs: vascular diseases.**

univariate model showed that multiple factors were significantly associated with poor outcomes (Table 4). However, in the multivariate model, only higher serum urea, high systolic blood pressure (SBP), and lower estimated glomerular filtration rate were associated with inferior outcomes. Severe glomerulosclerosis (GS) and interstitial fibrosis (IF) were significantly associated with inferior outcomes, with the highest adjusted odds ratio for severe IF (adjusted odds ratio [AOR], 6.41[2.19–18.76],  $p$  0.001).

## Discussion


In this study, we investigated the histopathological findings of renal biopsies in patients with unexplained renal impairment.

We found that glomerular diseases were the most common histological diagnosis and that severe GS and IF were significantly associated with inferior outcomes.


Every geographical area has a unique renal disease profile that becomes more evident with histopathological analysis. In our study, we found the most common histological diagnosis was glomerular diseases (41.9%). This was in agreement with Mittal et al., who found that 60% of patients presenting with CKD of undetermined etiology in Northern India had glomerular diseases.<sup>11</sup> This finding may reflect a delay in diagnosis as the majority of patients presented in advanced stages. Moreover, the median creatinine in our patients was 724 μmol/L, which was much higher than that reported by Syahril et al. (306 μmol/L) who investigated the renal histology in patients presenting by unexplained elevated serum creatinine in Malaysia. This reflects the problem of delayed referral to nephrologists in our region.<sup>12</sup>

The leading pathologies for glomerular diseases were FSGS (14.3%) and amyloidosis (11.4%). Similarly, in a study by Mittal et al., FSGS was the leading primary glomerulonephritis accounting for 18.2% of the cohort.<sup>11</sup> In contrast, Zaza et al. found that the leading types of glomerulonephritis detected from renal biopsies in Italian patients with CKD were IgA (24.3%) nephropathy and FSGS (13.3%).<sup>13</sup> Nevertheless, we did not recognize cases of IgA nephropathy among our patients. This may be explained by the lower prevalence of IgA in Africa.<sup>14</sup>

A definite cause for renal impairment could not be determined in 10.4% of the cohort. This was close to that reported in previous studies<sup>15,16</sup> and lower than reported by Titze et al.<sup>17</sup> Of them, 2.3% had clinical i.e., positive family history, positive consanguinity, or being young adults, and tubulointerstitial findings favoring a diagnosis of genetic origin i.e. tubular basement membrane thickening, lamellation, and multilayering



**Fig. 4 – Histopathological findings of the studied groups. CKDu: chronic kidney disease of unknown etiology, GS: glomerulosclerosis, IF: interstitial fibrosis, PRDs: paraprotein-related renal diseases, TA: tubular atrophy, TIDs: tubulointerstitial diseases, VDAs: vascular diseases. P1 comparison between glomerulosclerosis among whole groups by Chi-square test, P2 comparison between tubular atrophy among whole groups by Chi-square test, P3 comparison between interstitial fibrosis among whole groups by Chi-square test, P4 comparison between arterial sclerosis among whole groups by Chi-square test.**



**Fig. 5 – treatment strategies for studied groups. CKDu: chronic kidney disease of unknown etiology, PRDs: paraprotein-related renal diseases, TIDs: tubulointerstitial diseases, VDAs: vascular diseases.**

288 or presence of microcytes on EM examination. Genetic testing  
 289 to confirm the diagnosis was not done due to financial  
 290 constraints.

291 The other 8.1% had histopathological findings suggestive  
 292 of CKDu, such as non-specific glomerular sclerosis, chronic

interstitial nephritis, interstitial fibrosis, or lymphocytic  
 293 infiltration on LM, negative immunostaining and nonspecific  
 294 advanced chronic changes in EM. This was comparable to  
 295 (7.4%) that was reported by Foula et al.<sup>6</sup> This percentage is  
 296 more amenable than previously thought to decrease if genetic  
 297

**Table 3 – patient's outcome by histological group.**

Variables	All cohort, N = 210	CKDu, n = 17	GDS, n = 88	PRDs, n = 27	TIDs, n = 43	VDs, n = 35
Improved	43 (20.5%)	0 (0.0%)	7 (8.0%)	6 (22.2%)	26 (60.5%)	4 (11.4%)
CKD ND	76 (36.2%)	3 (17.6%)	37 (42%)	8 (29.6%)	9 (20.9%)	19 (54.3%)
CKD-D	88 (41.9%)	14 (82.4%)	44 (50.0%)	11 (40.7%)	8 (18.6%)	11 (31.4%)
Death	3 (1.4%)	0 (0.0%)	0 (0.0%)	2 (7.4%)	0 (0.0%)	1 (2.9%)

Values represent frequency (percentage).

CKD-D: chronic kidney diseases on dialysis, CKD-ND: chronic kidney diseases not on dialysis, CKDu: chronic kidney disease of unknown etiology, GDS: glomerular diseases, PRDs: paraprotein-related renal diseases, TIDs: tubulointerstitial diseases, VDs: vascular diseases.

**Table 4 – The variables associated with poor outcome on univariate and multivariate analysis.**

Predictors	Univariate OR (95% C.I.)	p-Value	Multivariate AOR (95% C.I.)	p-Value
Age	0.99 (0.97–1.01)	0.519		
Urea	1.05 (1.02–1.07)	<0.001	1.03 (1.01–1.06)	<b>0.025</b>
Creatinine	1.02 (1.001–1.03)	<0.001		
iPTH	1.03 (1.002–1.05)	<0.001		
eGFR EPI	0.88 (0.83–0.93)	<0.001	0.90 (0.84–0.97)	<b>0.006</b>
SBP	1.027 (1.01–1.05)	<b>0.007</b>	1.03 (1.01–1.05)	<b>0.019</b>
MAP	1.03 (1.01–1.06)	<b>0.039</b>		
24 h protein	1.00 (1.00–1.00)	0.097		
GS ≥ 50%	5.68 (2.79–11.56)	<0.001	2.63 (1.05–6.60)	<b>0.040</b>
IF ≥ 50%	10.19 (4.26–24.35)	<0.001	6.41 (2.19–18.76)	<b>0.001</b>
Marked TA	4.07 (1.93–8.53)	<0.001		
AS	2.48 (1.42–4.35)	<b>0.001</b>		

Model used was backward Wald logistic regression.

p value significant < 0.05 and shown in bold.

AOR: adjusted odds ratio, As: arterial sclerosis, CI: confidence interval, eGFR: estimated glomerular filtration rate, GS: glomerulosclerosis, IF: interstitial fibrosis, iPTH: intact parathyroid hormone, MAP: mean blood pressure, OR: odds ratio, SBP: systolic blood pressure, TA: tubular trophy.

testing was involved as concluded in recent studies that found 10–40% of cases of CKDu could be due to undiagnosed genetic diseases<sup>15,18,19</sup> or genetic predisposition to CKDu.<sup>20–22</sup>

All patients with findings suggestive of CKDu were inhabitants of rural areas, but only 47% reported working as farmers cultivating sugarcane, wheat, or vegetables. Except for two young patients, the other nine patients were middle-aged, close to the age reported in India and Sri Lanka, with male sex predominance.<sup>2,11</sup> None of them were diabetic, nor had a positive family history of renal diseases, or extrarenal manifestations. They had higher serum creatinine levels than reported in the literature. This may be due to the insidious onset and slow progression of their disease.

Amyloidosis (other than light chain amyloidosis) was found in 15% of our patients, which exceeded the percentage reported by Zaza et al. (7%).<sup>13</sup> This may reflect the prevalence of underrecognized amyloidosis among Egyptians, especially the ALECT2 phenotype.<sup>23</sup> Moreover, unsuspected light chain cast nephropathy was detected in 22 patients (10.5%). However, few cases had a similar presentation in the published literature.<sup>24,25</sup> This supports the importance of multiple myeloma screening in cases of unexplained renal injury.<sup>26</sup> We found that 12.6% of patients had vasculitis despite negative ANCA serology. This highlights the role of renal biopsies and not relying on serological tests to rule out this organ, or even life-threatening disease.

Forty percent of our patients had potentially treatable diagnoses by directed therapy beyond cardiovascular and CKD-MBD optimization. This was in agreement with Zaza et al.

who found that treatment strategies remained unchanged after biopsy in patients with CKD, and only symptomatic therapy was continued in 60% of patients.<sup>13</sup> However, Kitterer D et al. reported that renal histology resulted in identifying diseases potentially responsive to treatment modification in 74% of cases.<sup>27</sup> This discrepancy may be attributed to the fact that most of our patients presented with advanced stages, as well as the presence of diagnoses that do not have a definite standard of care treatment i.e., CKDu, some genetically inherited disorders, or LECT amyloidosis. Besides, treatment modifications were undergone after biopsy results such as discontinuation of immunosuppression for patients with secondary FSGS, searching for inflammatory conditions for patients with AA amyloidosis, and cessation of the offending factors in ATN cases.

## Conclusion

Our study underlines the importance of renal biopsy in patients with unexplained renal impairment. It has a fundamental role in disease characterization, prognostication, and treatment optimization even in the presence of the advanced nature of renal involvement. Moreover, transplant decisions such as the risk of recurrence of the original disease, the possibility of relative donation, and the need for simultaneous organ transplantation have a pertinent relation to renal biopsy findings. In patients with unexplained renal impairment, glomerular diseases were the most common his-



tological diagnosis, and FSGS and amyloidosis were the most frequent glomerular pathologies. Severe GS and IF were significantly associated with inferior outcomes. Finally, a more integrated primary healthcare program is needed to expedite early diagnosis, timely management, and subsequently prevent progression to CKD.

### Authors' contributions

Effat A. Tony: This author helped study design, writing, and editing of the manuscript.

Rabab Radi: This author helped data collection and excel sheets preparation.

Wesam Ismail: This author helped conduct of the study, data collection and draft writing of the manuscript.

Mohamed Ismail Seddik: This author helped conduct of the study and data collection.

Radwa A. Ellisy: This author helped the design of the study, data collection and draft writing of the manuscript.

Essam M. Abdel Aziz: This author helped in the final writing and editing of the manuscript.

### Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee at which the studies were conducted (IRB approval number IRB17200010), and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

### Funding

The authors have no sources of funding to declare for this manuscript.

### Conflicts of Interest

The authors have declared that no conflict of interest exists.

The authors declare no conflicts of interest

### Data sharing information

Raw data (de-identified) used in this clinical trial are available from the corresponding author. It will be available following publication upon reasonable request.

It has not been published in other journals in whole or part.

It was prospectively registered at [www.clinicaltrials.gov](http://www.clinicaltrials.gov) number; NCT03586531.

### REFERENCES

1. Foreman KJ, Marquez N, Dolgert A, Fukutaki K, Fullman N, McGaughey M, et al. Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes

of death: reference and alternative scenarios for 2016–40 for 195 countries and territories. *Lancet Lond Engl*. 2018;392:2052–90.

2. Johnson RJ, Wesseling C, Newman LS. Chronic kidney disease of unknown cause in agricultural communities. *N Engl J Med*. 2019;380:1843–52.

3. Weaver VM, Fadrowski JJ, Jaar BG. Global dimensions of chronic kidney disease of unknown etiology (CKDu): a modern era environmental and/or occupational nephropathy? *BMC Nephrol*. 2015;16:145.

4. El Minshawy O. End-stage renal disease in the El-Minia Governorate, upper Egypt: an epidemiological study. *Saudi J Kidney Dis Transplant*. 2011;22:1048–54.

5. Naiker IP, Assounga AG, Meyers AM. Diagnostic approach to chronic kidney disease. *S Afr Med J*. 2015;105:236.

6. Foula H, Zaghoul S, Elrewhby W, Abo Shosha H, Anwar M, Heddaya A, et al. Indications and histopathological patterns of kidney biopsies in Egypt: a single center, prospective analysis. *Kidney Int Rep*. 2020;5 Suppl.:S366–7.

7. Varghese RT, Jialal I. Diabetic nephropathy. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2023. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK534200/>

8. Clinical features, diagnosis, and treatment of hypertensive nephrosclerosis – UpToDate. Available from: <https://www.uptodate.com/contents/clinical-features-diagnosis-and-treatment-of-hypertensive-nephrosclerosis>.

9. Hogan JJ, Mocanu M, Berns JS. The native kidney biopsy: update and evidence for best practice. *Clin J Am Soc Nephrol*. 2016;11:354–62.

10. Pannu N, James M, Hemmelgarn B, Klarenbach S. Network for the AKD. Association between AKI, recovery of renal function, and long-term outcomes after hospital discharge. *Clin J Am Soc Nephrol*. 2013;8:194.

11. Mittal P, Agarwal SK, Singh G, Bhowmik D, Mahajan S, Dinda A, et al. Spectrum of biopsy-proven renal disease in northern India: a single-centre study. *Nephrol Carlton Vic*. 2020;25:55–62.

12. Ahmad Syahril WRWA, Chee Eng C, Fariz Safhan MN, Khadijah MN. Renal biopsy outcomes in patients with unexplained rise in serum creatinine. *Kidney Int Rep*. 2017;2:S22.

13. Zaza G, Bernich P, Lupo A. 'Triveneto' Register of Renal Biopsies (TVRRB). Renal biopsy in chronic kidney disease: lessons from a large Italian registry. *Am J Nephrol*. 2013;37:255–63. <http://dx.doi.org/10.1159/000348566>. Epub 2013 Mar 13. PMID: 23485642.

14. Storrar J, Chinnadurai R, Sinha S, Kalra PA. The epidemiology and evolution of IgA nephropathy over two decades: a single centre experience. *PLOS ONE*. 2022;17, e0268421.

15. Robert T, greillier S, Torrents J, Raymond L, Dancer M, Jourde-Chiche N, et al. Diagnosis of kidney diseases of unknown etiology through biopsy-genetic analysis. *Kidney Int Rep*. 2023;8:2077–87.

16. Robert T, Raymond L, Dancer M, Torrents J, Jourde-Chiche N, Burtsey S, et al. Beyond the kidney biopsy: genomic approach to undetermined kidney diseases. *Clin Kidney J*. 2023;17:sfad099.

17. Titze S, Schmid M, Köttgen A, Busch M, Floege J, Wanner C, et al. Disease burden and risk profile in referred patients with moderate chronic kidney disease: composition of the German Chronic Kidney Disease (GCKD) cohort. *Nephrol Dial Transplant*. 2015;30:441–51.

18. Robert T, greillier S, Torrents J, Raymond L, Dancer M, Jourde-Chiche N, et al. Diagnosis of kidney diseases of unknown etiology through biopsy-genetic analysis. *Kidney Int Rep*. 2023;8:2077–87.

19. Groopman EE, Marasa M, Cameron-Christie S, Petrovski S, Aggarwal VS, Milo-Rasouly H, et al. Diagnostic utility of

- 457 exome sequencing for kidney disease. *N Engl J Med.* 2019;380:142–51. 476
- 458 20. Nanayakkara S, Senevirathna STMLD, Abeysekera T, 477
- 459 Chandrajith R, Ratnatunga N, Gunarathne EDL, et al. An 478
- 460 integrative study of the genetic, social and environmental 479
- 461 determinants of chronic kidney disease characterized by 480
- 462 tubulointerstitial damages in the North Central Region of Sri 481
- 463 Lanka. *J Occup Health.* 2014;56:28–38. 482
- 464 21. Nanayakkara S, Senevirathna STMLD, Parahitiyawa NB, 483
- 465 Abeysekera T, Chandrajith R, Ratnatunga N, et al. 484
- 466 Whole-exome sequencing reveals genetic variants associated 485
- 467 with chronic kidney disease characterized by 486
- 468 tubulointerstitial damages in North Central Region, Sri Lanka. 487
- 469 *Environ Health Prev Med.* 2015;20:354–9. 488
- 470 22. Siddarth M, Datta SK, Ahmed RS, Banerjee BD, Kalra OP, 489
- 471 Tripathi AK. Association of CYP1A1 gene polymorphism with 490
- 472 chronic kidney disease: a case–control study. *Environ Toxicol* 491
- 473 *Pharmacol.* 2013;36:164–70, 492
- 474 <http://dx.doi.org/10.1016/j.etap.2013.03.008>. Epub 2013 Mar 21. 493
- 475 PMID: 23619522. 494
23. Larsen CP, Ismail W, Kurtin PJ, Vrana JA, Dasari S, Nasr SH. 476
- Leukocyte chemotactic factor 2 amyloidosis (ALECT2) is a 477
- common form of renal amyloidosis among Egyptians. *Mod* 478
- Pathol.* 2016;29:416–20. 479
24. Adachi G, Mouri N, Ohta R, Sano C. A case of cast 480
- nephropathy found as the cause of severe renal failure. 481
- Cureus.* 2021;13:e19135. 482
25. Gastelum ZN, Biggs DM, Scott A. Multiple myeloma 483
- presenting as acute renal failure in the absence of other 484
- characteristic features. *Cureus.* 2017;9:e1703, 485
- <http://dx.doi.org/10.7759/cureus.1703>. PMID: 29159010; 486
- PMCID: PMC5690288. 487
26. Yadav P, Sathick IJ, Leung N, Brown EE, Cook M, Sanders PW, 488
- et al. Serum free light chain level at diagnosis in myeloma 489
- cast nephropathy—a multicentre study. *Blood Cancer J.* 490
- 2020;10:28. 491
27. Kitterer D, Gürzing K, Segerer S, Alscher MD, Amann K, Braun 492
- N, et al. Diagnostic impact of percutaneous renal biopsy. *Clin* 493
- Nephrol.* 2015;84:311–22. 494