Healthcare models, quality indicators and quality of care in the management of patients with chronic kidney disease in Spain: results from the CARABELA-CKD initiative

Mercedes Salgueira Pedro Ruiz Jesús Ignacio Diago Manuel Pérez Maraver, CARABELA-CKD Scientific Committee



PII: S0211-6995(24)00110-3

DOI: https://doi.org/doi:10.1016/j.nefro.2024.11.001

Reference: NEFRO 1299

To appear in: NEFROLOGÍA

Received Date: 14 August 2024

Accepted Date: 12 November 2024

Please cite this article as: Salgueira M, Ruiz P, Diago JI, Maraver MP, Healthcare models, quality indicators and quality of care in the management of patients with chronic kidney disease in Spain: results from the CARABELA-CKD initiative (2024), doi: https://doi.org/10.1016/j.nefro.2024.11.001

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U.

Healthcare models, quality indicators and quality of care in the management of patients with chronic kidney disease in Spain: results from the CARABELA-CKD initiative

Modelos asistenciales en salud, indicadores de calidad y calidad de la atención en el manejo de los pacientes con enfermedad renal crónica en España: resultados de la iniciativa CARABELA-ERC

Authors: Mercedes Salgueira^{a, b*}, Pedro Ruiz^{c, d*}, Jesús Ignacio Diago^e, CARABELA-CKD Scientific Committee[#] and Manuel Pérez Maraver^{f, g&}

Institutions:

- 1. Spanish Society of Nephrology (S.E.N.)
- 2. Nephrology Department. University Hospital Virgen Macarena, Seville, Spain
- 3. Spanish Society for Healthcare Quality (SECA)
- 4. AMCA Asociación Madrileña de Calidad Asistencial, Madrid, Spain
- 5. AstraZeneca Medical Department, Madrid, Spain
- 6. Spanish Society of Endocrinology and Nutrition (SEEN)
- 7. Endocrinology Department, Bellvitge University Hospital, Barcelona, Spain

*List and affiliation of the CARABELA-CKD Scientific Committee members are shown in the appendix

&Corresponding author

Manuel Pérez Maraver

Spanish Society of Endocrinology and Nutrition (SEEN); Endocrinology Department, Bellvitge University Hospital, Barcelona, Spain.

Carrer de la Feixa Llarga, s/n, 08907 L'Hospitalet de Llobregat, Barcelona mmperez@bellvitgehospital.cat

^{*}Both authors contributed equally to this manuscript

Healthcare models, quality indicators and quality of care in the management of patients with chronic kidney disease in Spain: results from the CARABELA-CKD initiative

Modelos asistenciales en salud, indicadores de calidad y calidad de la atención en el manejo de los pacientes con enfermedad renal crónica en España: resultados de la iniciativa CARABELA-ERC

ABSTRACT

Background and objectives: Chronic kidney disease (CKD) poses a significant public health challenge worldwide and its prevalence is rising due to aging populations and increasing risk factors. The CARABELA-CKD initiative aims to address CKD management inefficiencies. Materials and methods: The CARABELA-CKD initiative was structured into four phases. The methodology involved five pilot center studies, conducted to characterize CKD healthcare models (Phase 1), followed by the validation of disease care models, improvement areas, potential solutions, and strategic healthcare quality indicators in a National Multidisciplinary Workshop (Phase 2). Regional meetings were then held to refine all these aspects from a regional perspective and to determine the cocreation of solutions with subsequent quantitative validation of results using the two-round Delphi method (Phase 3B). Dissemination and implementation included integrating findings into digital questionnaires creating digital Playbooks and organizing local workshops (Phase 4). Results: The study identified three CKD care models in Spanish hospitals, each with distinct levels of specialization and resource availability. Twenty-five healthcare quality indicators were validated in a national meeting to assess various aspects of care quality and transformation. Improvement areas were identified, leading to the formulation of solutions to enhance CKD care models in Spain. Conclusions: The CARABELA-CKD initiative, a collaborative effort between Scientific Societies involved in the CKD care process and AstraZeneca, is a nationwide attempt in Spain to create a

framework to promote solutions in healthcare delivery and improvements in the quality of care that result in better health outcomes and higher standards of care quality, improved quality of life for patients, and increased efficiency of the Healthcare Service.

Keywords: chronic kidney disease, healthcare models, indicators, quality of care, management

RESUMEN

Antecedentes y objetivo: La enfermedad renal crónica (ERC) representa un desafío significativo para la salud pública a nivel mundial. Su prevalencia está en aumento debido al envejecimiento poblacional y a factores de riesgo crecientes. La iniciativa CARABELA-CKD busca abordar las ineficiencias en el manejo de la ERC. Materiales y métodos: La iniciativa CARABELA-CKD se estructuró en cuatro fases. En la primera fase, se caracterizaron los modelos de atención de la ERC en cinco centros piloto. En una segunda fase en una Reunión Nacional se validaron estos modelos, las áreas de mejora y las soluciones potenciales, además de indicadores de calidad en el manejo de la ERC. En la tercera fase se procedió a realizar reuniones regionales que facilitaron el refinamiento de las soluciones bajo una perspectiva regional con validación cuantitativa de resultados mediante un proceso Delphi (Fase 3B). En la cuarta fase se integraron los hallazgos en cuestionarios digitales permitiendo, mediante talleres, la diseminación e implementación local de CARABELA-CKD. Resultados: El estudio identificó tres modelos de atención de la ERC en hospitales españoles, cada uno con niveles distintos de especialización y disponibilidad de recursos. Se validaron 25 indicadores para evaluar varios aspectos de la calidad de atención y transformación. Se identificaron áreas de mejora, lo que llevó a la formulación de soluciones para mejorar los modelos de atención de la ERC en España. Conclusiones: La iniciativa CARABELA, entre las sociedades científicas relacionadas con el manejo de la ERC y AstraZeneca, es una iniciativa multidisciplinar a nivel nacional de crear un marco para promover cambios en la atención sanitaria que resulten en

mejoras en los resultados en Salud y en estándares más altos de calidad asistencial, mejora en calidad de vida de los pacientes y eficiencia del sistema sanitario.

Palabras clave: enfermedad renal crónica, modelos asistenciales, indicadores, calidad asistencial, manejo

INTRODUCTION

Chronic kidney disease (CKD) is a public health problem worldwide and is considered a silent epidemic due to the low rate of diagnosis and the lack of awareness of this condition among the population ¹⁻⁶. Approximately 15% of the Spanish population is affected by CKD ⁴. The progressive aging of Western society, coupled with the increasing prevalence of arterial hypertension, diabetes mellitus, obesity and heart failure are crucial factors that are driving the progressive rise in the prevalence of CKD. ^{1,7}. This results in a heightened risk of premature mortality, especially from cardiovascular causes. If the current trend continues, by 2040, CKD will be the fifth leading cause of death in Spain and by the end of the century, it may possibly rank as the second leading cause of death in Spain ⁶. Furthermore, there has been a 30% overall rise in advanced CKD necessitating kidney replacement therapies in recent decades in Spain, impacting over 64,000 individuals. The economic burden is substantial, accounting for 3% of total healthcare costs ⁸. Therefore, developing and implementing strategies aimed at optimizing the management of CKD is absolutely necessary.

The CARABELA-CKD is a strategic initiative developed within the CARABELA framework, designed to tackle the intricate improvement areas associated with CKD management within the Spanish Healthcare Service ⁹. CARABELA initiatives aim to achieve a thorough understanding of the improvement areas that all stakeholders involved in the care of chronic diseases in the Healthcare Service must address in order to catalyze the transformation in

clinical practice needed to achieve improvements in the quality of care. CARABELA initiatives are rooted in lean process reengineering and constitute holistic efforts in which Scientific Societies and AstraZeneca collaborate to examine current improvement areas in chronic diseases. In sum, CARABELA-CKD, spearheaded by a multidisciplinary committee, with the participation of 3 Scientific Societies, represents a comprehensive and transversal initiative aimed at addressing management inequalities and identifying inefficiencies in CKD management in Spain.

This article presents the outcomes of the CARABELA-CKD initiative which focus on assessing the existing CKD management models in Spain to identify improvement areas, solutions and healthcare quality indicators to delineate the action plans.

MATERIALS AND METHODS

Design and participants of CARABELA-CKD

The overall structure and basis of the CARABELA methodology have been documented previously ⁹. Specifically, CARABELA-CKD comprised four phases (Figure 1). Coordination throughout Spain was carried out by the Scientific Societies involved in the care of patients with CKD, namely S.E.N. (Spanish Society of Nephrology), SEEN (Spanish Society of Endocrinology and Nutrition), and SECA (Spanish Society for Healthcare Quality), and AstraZeneca.

Characterization of CKD healthcare models in Spanish hospitals (Phase 1)

The characterization phase was framed on the understanding that CKD care is a circular and multidisciplinary process ¹⁰. To start, a general model or patient pathway for CKD clinical management was developed by the Scientific Committee and tested in five pilot centers representing the Spanish geography and organizational models, in order to identify variations and adaptations to different settings (Figure 1). This process led to the categorization of various models of patient care. To evaluate each of the identified models and serve as reference for future monitoring, improvement areas, solutions and a list of healthcare quality indicators were developed by the Scientific Committee for CARABELA-CKD. Results from the first phase were validated by the Scientific Committee in preparation for the National Conference.

Validation of results (Phase 2) and regional (Phase 3) and local (Phase 4) phases.

A National Multidisciplinary Workshop was held with 100 representatives from the CKD healthcare ecosystem in Spain. In this meeting, the following actions were implemented: validation of the nine variables that would define and characterize each care model, selection of the strategic healthcare quality indicators using a two-round Delphi survey, and validation of improvement areas and potential solutions applicable to the different CKD healthcare models. The results of the National Multidisciplinary Workshop were further shared and discussed in a set of four Regional Multidisciplinary Meetings with the participation of physicians representative of the whole Spanish territory, in which local and regional realities were further considered, providing additional insight and opportunities to propose regionally refined solutions and adaptations for various settings. Based on the findings and discussions from the National Multidisciplinary Workshop and the Regional Multidisciplinary Meetings within the CARABELA-CKD initiative, solutions were formulated to address improvement areas within the CKD care process. At present, CARABELA-CKD results serve as the basis for an ongoing practical phase, in which improvement areas and solutions are implemented into digital questionnaires in the form of Playbooks. Local workshops to analyze the model of healthcare units and design tailored actions toward improvement in the care of CKD are organized in healthcare areas throughout Spain upon request.

RESULTS

Coordination models for the management of CKD in Spanish hospitals

During the pilot phase conducted in five centers, three CKD patient care models were identified based on nine variables used for characterizing the different models of approach (Figure 2). These variables were associated with the type and level of coordination among the medical specialties involved in CKD care, the roles of the professionals involved, and the

availability of specific resources. A total of nine variables were identified and analyzed for the characterization of each model: 1) coordination model or multidisciplinary approach among specialties integrating nephrology, endocrinology and/or cardiology; 2) availability of skilled CKD nursing staff; 3) availability of specialized nursing care with dedicated schedules; 4) availability of advanced multidisciplinary CKD units; 5) presence of specialized nursing staff, trained and dedicated to patient care in multidisciplinary CKD units; 6) coordination, referral protocols and channels of communications among levels of care; 7) availability of case managers for managing hospital transitions, admissions and discharges); 8) availability of a multipurpose day hospital area with high specialization in CKD; and 9) availability of tools to assess patient outcomes (Patient-Reported Experience Measures [PREMs] and Patient-Reported Outcomes Measures [PROMs]).

Among three CKD patient care models, the main differentiating factor was the availability of multidisciplinary CKD units and specialized CKD nursing. Diversity and accessibility of resources for addressing CKD were differentiating factors among the models (Figure 2). Additionally, a detailed description of the patient journey according to the phase of the care process (diagnosis, treatment, and follow-up) were described in Supplementary Material.

In Model 1, CKD units, multidisciplinary units are in place for cardiorenal and diabetic nephropathy care (renovascular unit optional) provided by specialized and trained nursing staff working in multidisciplinary units for CKD care that are adequately equipped with the necessary resources.

In **Model 2, CKD units** are characterized by the presence of at least one of the three multidisciplinary units mentioned: cardiorenal, diabetic nephropathy, or renovascular care. This model has a nephrology-specific nursing staff, exclusively dedicated to the care of patients with CKD (at any stage, including advanced CKD treated in multidisciplinary units). It is important to note that the nursing staff has a separate physical space and a dedicated schedule to attend to CKD patients.

Model 3 is distinguished by the fact that the nursing staff who care for CKD patients lack specific specialization, and only exceptionally have their own physical space and dedicated schedules.

Regarding communication between levels of healthcare and among specialties, models 1 and 2 differed in that the first proposes a multidisciplinary approach with bidirectional communication channels, while the second proposes "coordination between specialties".

Furthermore, Model 1 includes a multipurpose day hospital exclusively and specifically for the use of the Nephrology Unit, while in Model 2, the general hospital day unit is used. Lastly, Model 1 suggested the occasional presence of a case manager and the collection of PROMS and PREMS, features that were not available in either Model 2 or Model 3.

Healthcare quality indicators for the evaluation of CKD care models

The proposed indicators for evaluating the care provided in the CKD process in each unit include structural, quality of care, and transformation indicators (Table 1) (detailed in the materials and methods).

A total of 156 potential indicators were initially proposed by the members of the Scientific Committee. These subsequently underwent a voting process to prioritize and select the most appropriate, and finally, the selected indicators were shared and voted on using Delphi methodology at the National Multidisciplinary Workshop and further validated in regional meetings and by the Scientific Committees. The selection criteria were feasibility of implementation and potential beneficial impact for improving patient care or healthcare results. Twenty-five indicators were selected in a two-round Delphi process at the National Workshop by 100 clinical experts. The selected indicators were presented to participants at the four Regional Multidisciplinary Meetings, in which healthcare professionals from across the

country provided their feedback. The final list was confirmed by the Scientific Committee (Table 1).

The category of structural indicators focuses on how the center is organized for the management of CKD (Table 1). Healthcare quality indicators were divided according to the phase of the CKD care process: diagnosis, treatment, and follow-up (Table 1). Likewise, the Scientific Committees defined different types of transformation indicators that would offer a comprehensive view of the evolution of the care model: transformation of the structure, transformation of the process, and improvement areas.

Future improvement areas for the evolution of CKD care models: barriers and potential solutions

The comprehensive analysis identified improvement areas, which were validated in the CARABELA-CKD National Multidisciplinary Workshop. They were grouped as four major challenges: care process, social, technological, and organizational and economic solutions, all of which will be addressed in the future to establish an integrated care model.

These improvement areas were also discussed and validated at the Regional Multidisciplinary Meetings with healthcare professionals, identifying solutions (grouped as eight axes of change and 17 lines of action) (Table 2). The professional analysis and expert opinions that led to this list will be the topic of a dedicated publication.

DISCUSSION

CKD is a complex condition often overlooked in its early stages ⁶. Hence, there is an urgent need to design a coordinated strategy aimed at improving the management of individuals with CKD to implement the most effective approach for reducing mortality and healthcare expenses ¹¹. Effective management should include addressing CKD risk factors ¹⁰ and associated conditions and comorbidities ¹² to prevent disease progression. Indeed, a few years ago, the

Commitment to Quality project led by the Spanish Society of Nephrology (S.E.N.), in coordination with the Ministry of Health, Consumption and Social Welfare, set four objectives and solutions aimed at addressing CKD comprehensively. These solutions focus on early-stage disease identification and commit to delivering high-quality, patient-centered care throughout the patient journey, starting from the initial symptoms and encompassing all stages of CKD and clinical progression ^{7,8,11,13}.

Nonetheless, the current state of the care process in our country is exceptionally diverse, reflecting a range of scenarios in the management of CKD patients. This diversity is influenced by factors such as geographical location, cultural considerations, and the distinctive characteristics of the various regional health systems ⁹. Consequently, it has become imperative to assess the implementation of quality-of-care programs and gain a comprehensive understanding of the current management of CKD in Spanish healthcare centers. Hence, CARABELA-CKD has concentrated its efforts on not only identifying models, but also assessing resource utilization and management characteristics. Only through this approach can we identify limitations, transform the system toward adopting effective and efficient organization and approaches, and develop targeted solutions that may be either traditional or innovative.

The results from the CARABELA-CKD in-depth characterization process conducted in five pilot centers shed light on the distinct scenarios and on the diverse levels of accessibility to resources within the management of CKD. This research has identified three care models for patients with CKD in Spanish hospitals categorized according to specialization, availability of resources, multidisciplinary care, and specialized nursing. In these models, the most significant and impactful factor is multidisciplinary collaboration, embodied by functional units, adequately equipped with the necessary resources for the management of CKD. Nonetheless, each of these models has the potential for improvement and the attainment of excellence. Additionally, a diverse set of indicators, encompassing structural, quality-of-care and

transformation indicators were collected and validated to help enhance the CKD care process. These indicators provide valuable insights into critical aspects of care, enabling ongoing monitoring and guiding continuous improvement efforts ¹². They are tailored to each of the identified care models and processes, thus facilitating improvements within each model that align with its distinct realities, requirements, and available resources. These improvements include effective communication and coordination among all healthcare professionals managing CKD, the promotion of education and training for both healthcare professionals and the patients themselves, and the use of protocols that incorporate referral criteria to standardize CKD care. In this regard, we draw upon previous experience from other countries, where the implementation of integrated programs has yielded notable enhancements and improvements in the care of various clinical entities ¹⁴⁻¹⁶.

When compared with other strategic initiatives, CARABELA-CKD is an innovative proposition that takes into account the diverse realities and existing care models across Spain ⁹. It envisions the care of CKD patients in Spain as a circular and multidisciplinary process, involving the seamless integration of hospital care areas, such as nephrology, endocrinology, cardiology, and primary care, all operating cohesively within a patient-centered framework. This is due to the ability of the initiative to furnish a comprehensive, holistic and pragmatic perspective across various settings and scenarios, along with a tailored improvement framework that adapts to diverse contexts. Moreover, CARABELA-CKD represents a collaborative initiative that brings together both Scientific Societies and AstraZeneca, uniting clinical leadership with an integrative philosophy.

Throughout the development of this initiative, participants have evaluated, deliberated, and critically scrutinized care models and indicators, identified improvement areas, and proposed potential solutions. The participation of Scientific Societies from different specialties and healthcare professionals directly involved in the care of patients with CKD, both at the national and regional levels, adds value to this initiative.

Finally, in a subsequent practical phase, the dissemination and implementation of this initiative in the form of local workshops held in healthcare centers throughout the country will facilitate its implementation and bring about a transformation in the current care model for renal patients.

Since the CARBELA-CKD provides a robust methodology that enables each healthcare center or team to position itself within each specific model, it offers the potential for enhancement while aligning meticulously with the demands of real-world healthcare processes. In essence, each model can discern its unique reference point for improvement. Thus, through self-analysis, they can craft their own improvement plans, tailored to their individual circumstances. This transformation aims to optimize clinical pathways and the overall management of patients, ultimately leading to improved healthcare outcomes and reducing the burden of CKD.

CONCLUSIONS

In conclusion, the CARABELA-CKD initiative is a collaborative effort, involving different Scientific Societies, healthcare professionals across Spain, and AstraZeneca. CARABELA-CKD has defined a variety of existing CKD care models which can be grouped into three main models. It has also-identified strategic healthcare quality indicators in the renal patient care pathway, and has established improvement areas, applicable to all care models, and actions to be taken to enhance the quality of care for CKD. Its implementation may result in overall improvement in patient health outcomes and efficiency throughout the Healthcare Service.

Funding

All support for the present manuscript was provided by AstraZeneca Farmacéutica Spain.

Conflicts of interest

MS has received payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from Baxter, GSK, Boehringer, Astellas, AstraZeneca, and Vifor; has received support for attending meetings and/or travel from Vifor and Menarini; and has receipt of equipment, materials, drugs, medical writing, gifts or other services from Vifor and GSK. PR declares no conflict of interest. JID is an employee at the Medical Department of AstraZeneca Farmacéutica Spain. MPM has received grants or contracts from Sanofi; has received payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from AstraZeneca, Novo Nordisk, Lilly, Sanofi, Abbott, Boehringer, Bayer, and Menarini.

Acknowledgments

The authors acknowledge the participation of all the professionals attending the regional meetings of the CARABELA-CKD initiative and all those involved in the pilot phase, from Hospital Clínico Universitario de Valencia (Valencia, Spain), Hospital Universitario de Cruces (Vizcaya, Spain), Hospital Universitario 12 de Octubre (Madrid, Spain), Hospital Universitario Reina Sofía (Córdoba, Spain), Hospital Universitario Vall d'Hebron (Barcelona, Spain), Hospital Universitario Lucus Augusti (Lugo, Spain), Complejo Hospitalario Universitario de A Coruña (A Coruña, Spain), Hospital Universitario Marqués de Valdecilla (Santander, Spain), Hospital Universitario de Bellvitge (Barcelona, Spain), Hospital Universitario Juan XXIII (Tarragona, Spain), Hospital Universitario Doctor Josep Trueta (Girona, Spain), Hospital Universitario Fundación Jiménez Díaz (Madrid, Spain), Centro de Salud Rafael Alberti (Madrid, Spain), Hospital Universitario Virgen de las Nieves (Granada, Spain), Hospital Universitario Virgen Macarena (Seville, Spain), Hospital Universitario de Badajoz (Badajoz, Spain).

Medical writing support under the guidance of the authors was provided by Laura Vilorio Marqués, PhD, Blanca Piedrafita, PhD, and Javier Arranz-Nicolás, PhD, from Medical Statistics Consulting (MSC), Valencia, Spain, in accordance with Good Publication Practice guidelines (DeTora, L. M. et al. Ann Intern Med. 2022).

Appendix

The CARABELA-CKD Scientific Committee consists of the following members: Borja Quiroga (Sociedad Española de Nefrología, S.E.N.), Javier Escalada (Sociedad Española de Endocrinología y Nutrición, SEEN), Juan José Gorgojo (SEEN), Manuel Pérez Maraver (SEEN), Mercedes Salgueira (S.E.N.), Patricia de Sequera (S.E.N.), Pedro Ruiz (Sociedad Española de Calidad Asistencial), Alberto Prado Dominguez (AstraZeneca Farmacéutica Spain), Jesús Ignacio Diago (AstraZeneca Farmacéutica Spain), and Lucía Regadera (AstraZeneca Farmacéutica Spain).

References

- 1. Nefrología SEd. La Enfermedad Renal Crónica en España. In; 2022. Available at: https://www.seden.org/files/courses/Informe_390a.pdf [
- 2. Cobo Marcos M, de la Espriella R, Gayan Ordas J, Llacer P, Pomares A, Fort A, et al. Prevalence and clinical profile of kidney disease in patients with chronic heart failure. Insights from the Spanish cardiorenal registry. Rev Esp Cardiol (Engl Ed). 2023.
- 3. Exposito C, Pera G, Rodriguez L, Arteaga I, Martinez A, Aluma A, et al. Prevalence of Early Chronic Kidney Disease and Main Associated Factors in Spanish Population: Populational Study. J Clin Med. 2019;8(9).
- 4. Gorostidi M, Sanchez-Martinez M, Ruilope LM, Graciani A, de la Cruz JJ, Santamaria R, et al. Chronic kidney disease in Spain: Prevalence and impact of accumulation of cardiovascular risk factors. Nefrologia (Engl Ed). 2018;38(6):606-15.
- 5. iSanidad. La prevalencia de la enfermedad renal crónica sigue aumentando en España 2020 [Available from: https://isanidad.com/173405/la-prevalencia-de-la-enfermedad-renal-cronica-sigue-
- aumentando/#:~:text=La%20Enfermedad%20Renal%20Cr%C3%B3nica%20(ERC,tratamiento%20de%20di%C3%A1lisis%20o%20trasplante..
- 6. Airg E, Ekpf, Alcer, Friat, Redinren, Ricors, et al. CKD: The burden of disease invisible to research funders. Nefrologia (Engl Ed). 2022;42(1):65-84.
- 7. Prieto-Velasco M, Del Pino YPMD, Buades Fuster JM, Craver Hospital L, Pons Prades R, Ruiz San Millan JC, et al. Advanced Chronic Kidney Disease Units in Spain: a national survey on standards of structure, resources, results and patient safety. Nefrologia (Engl Ed). 2020;40(6):608-22.
- 8. Documento Marco sobre Enfermedad Renal Crónica (ERC) dentro de la Estrategia de Abordaje a la Cronicidad en el SNS (2015). Available at https://www.sanidad.gob.es/organizacion/sns/planCalidadSNS/pdf/Enfermedad_Renal_Cronic a_2015.pdf. In [
- 9. Escalada J, Carretero Gómez J, Anguita M, de Sequera P, García-Río F, Dávila I, et al. Enhancing the management of chronic diseases in clinical practices: The CARABELA methodology. Journal of Healthcare Quality Research. 2024;In press.
- 10. Quiroga B, Gorgojo JJ, Diago JI, Ruiz P. Toward optimization and excellence via a multidisciplinary care model for patients with chronic kidney disease in Spain: The CARABELA-CKD initiative. Nefrología.
- 11. Prieto-Velasco M, Aguilera Flórez AI, Agúndez Gómez I, Alaguero del Pozo B, Andrés Fernández B, Algarabel Cámara AA, et al. Proceso asistencial integrado para la prevención y la atención a las personas con enfermedad renal crónica 2018. Available from: https://www.saludcastillayleon.es/profesionales/es/procesos-asistenciales/procesos-asistenciales-gerencia-regional-salud/enfermedad-renal-cronica.ficheros/1367294-PAI.
- 12. Álvarez de Lara MA, Arenas MD, Salgueira Lazo M. Calidad y Seguridad en el Tratamiento del Paciente con Enfermedad Renal Crónica. Nefrología al día2021. Available from: https://www.nefrologiaaldia.org/es-articulo-calidad-seguridad-el-tratamiento-del-380.
- 13. MSSSI. Estrategia de abordaje de la Cronicidad http://www.msssi.gob.es/organizacion/sns/planCalidadSNS/pdf/ESTRATEGIA_ABORDAJE_CRO NICIDAD.pdf [
- 14. Goldstein SL. Automated/integrated real-time clinical decision support in acute kidney injury. Curr Opin Crit Care. 2015;21(6):485-9.
- 15. Wu HY, Fukuma S, Shimizu S, Norton EC, Tu YK, Hung KY, et al. Effects of Higher Quality of Care on Initiation of Long-term Dialysis in Patients With CKD and Diabetes. Am J Kidney Dis. 2017;70(5):666-74.

16. Heijsters F, van Breda FGF, van Nassau F, van der Steen MKJ, Ter Wee PM, Mullender MG, et al. A pragmatic approach for implementation of value-based healthcare in Amsterdam UMC, the Netherlands. BMC Health Serv Res. 2022;22(1):550.

Table 1. Structural, quality-of-care and transformation indicators validated in the CARABELA-CKD initiative according to type

AREA	TYPE OF	INDICATOR
	INDICATOR	
Human resources for CKD care	Structural	 Number of nephrologists in the department specialized in the care of advanced CKD (1/200 patients) Number of endocrinologists in the department dedicated to diabetes/10,000 habitants
		3) Number of nephrology nurses dedicated to the management of CKD / (1/200 patients)
The CKD patient pathway	Quality of care	 4) Percentage of patients with a waiting time for the first appointment of less than 30 days/Number of patients seen for the first time 5) Number of CKD patients admitted for hyperkalemia (>5.5 mmol/I)/Number of CKD patients) x100* 6) Number of correct referrals for CKD to the Nephrology unit according to local or national guidelines/Number of patients referred to the Nephrology clinic for suspected CKD) x100*. 7) Number of CKD patients with BMI ≥35 referred from Nephrology to Endocrinology
Care process	Quality of care	Diagnosis and treatment
Cure process	Quanty of care	8) Percentage of patients with glomerular filtration (GFR) estimation by Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) 9) Percentage of type 2 diabetic patients with a glomerular filtration rate greater than 60 ml/min in whom annual albuminuria testing is requested 10) Percentage of patients with albuminuria undergoing treatment with RAASi (ACEi/ARA-II) 11) Percentage of type 2 diabetic patients with CKD undergoing treatment with SGLT2i 12) Percentage of patients with HbA1c<7% out of the total number of patients analyzed 13) Percentage of patients with LDL<70 out of the total number of patients analyzed 14) Percentage of patients with blood pressure <130/80 out of the total number of patients in whom blood pressure is measured 15) Implementation of telematic
		communication of telematic communication systems between levels of care (Yes/No)

		16) Percentage of patients with cardiorenal metabolic syndrome referred to cardiorenal
		and/or diabetic nephropathy unit
	Quality of Care	Outcome indicators
		17) Percentage of patients requiring RRT
		annually out of total of CKD patients
		18) Existence of multidisciplinary clinical units
		of cardiorenal medicine
		19) Existence of multidisciplinary diabetic
		nephropathy clinical unit
		20) Percentage of patients in nephrology
		follow-up admitted to the hospital due to renal
		complications annually
		21) Percentage of patients on RRT who suffer a cardiovascular event annually
		22) Percentage of patients with CKD (at any
		stage) who die annually
		stage, who are annually
Transformation of	Transformation	23) Existence of a nephrology day hospital
structure, process, and		24) Existence of updated training support
improvement areas		materials for patients with CKD or caregivers
		validated by the center or Scientific Societies
		25) Existence of a local protocol aimed at the
		comprehensive management of patients with
		CKD (multidisciplinary assessment)

ACEI/ARA-II: angiotensin converting enzyme inhibitors/angiotensin II receptor antagonists; ACKD: advanced chronic kidney disease; BP: blood pressure; CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration; SGLT2i: sodium-glucose cotransporter-2 inhibitors; LDL: low-density lipoprotein; RAASi: Renin-angiotensin-aldosterone system inhibitors; RRT: renal replacement therapy.

Table 2: Potential solutions identified in the CARABELA-CKD initiative


AXE OF CHANGE	LINE OF ACTION
Coordination among healthcare professionals	1) Implementation of a case manager for hospital transitions, admissions and discharges 2) Implementation of multidisciplinary consultations that integrate diabetic nephropathy, cardiorenal and/or renovascular areas
Patient and professional training	3) Definition of training sessions conducted by nephrologists for all professionals involved in the management of CKD. 4) Creation and establishment of gamified health education programs (lifestyle and self-care)
Multidisciplinary protocols	 5) Existence of referral protocols to nephrology or day hospital from other units (emergency, PC, and other hospital specialties) 6) Existence of protocolized care pathways for chronic patients, with the participation of multidisciplinary groups
Advanced practice nurse in CKD	7) Existence of advanced CKD care units with dedicated and specialized nursing staff
Structured multidisciplinary care	8) Integration of nutrition, psychology, and occupational therapy specialties in the care and management of CKD patients
Promotion of a data-centric culture	9) Promotion of data recording through monitoring of essential indicators for the management of CKD 10) Development of training, educational and awareness programs for healthcare professionals for the implementation of comprehensive coding of patient pathologies
Resources to improve accessibility	11) Promotion of the growth of advanced CKD units 12) Creation of a multipurpose day hospital within nephrology units 13) Implementation of a streamlined communication channels between different care levels using information systems within the medical record 14) Promotion of home care for patients to prevent hospitalizations
Enhancement of patient experience and health outcomes	15) Standardization and development of tools to integrate patient-reported outcomes (PROMs and PREMs) into clinical practice

16) Development of health promotion and implementation
of general health and kidney prevention programs
17) Fostering patient empowerment and creation of
mentoring programs by initiating patient training in primary
care

CKD: chronic kidney disease; PC: primary care; PREM: patient-reported experiences; PROM: patient-reported outcomes; RRT: renal replacement therapy.

Figure 1: Flow chart of the CARABELA methodology. State of the art of the disease &

opportunities to optimize care



Phase 1: Characterization. 5 Pilot center study on the patient journey base and research of the CKD Healthcare models Collection and evaluation of key indicators in the 5 pilot centers. Identification of the different models of care, lines of action, and axes of change. Collection of information related to the patient experience.

Phase 2: Validation. First discussion towards validation and prioritization of 85 indicators by clinical and managing experts appointed by Scientific Societies in a National Meeting. Agreement on the models, the axes of change and the crosscutting lines of action identified, as well as the identification of 25 quality indicators.

Phase 3: Cocreation. In 4 regional meetings held with representatives of the regional healthcare ecosystem, the approach models identified were presented. Discussion and prioritization of the lines of action, identification of challenges and co-creation of solutions for their implementation follow.

Phase 3B: Quantitative validation. All previous Results and 25 Quality indicators were validated by the expert committees, by means of ad-hoc poll or Delphi consensus process.

Phase 4: Dissemination and Implementation. Results are implemented into digital questionnaires in the form of playbooks. Local workshops are organized upon request to analyze the model of healthcare units and design actions towards improvement.

Figure 2: Identified models, detailed according to the nine CKD care variables. CKD: chronic kidney disease; NA: not applicable; PREM: patient-reported experiences; PROM: patient-reported outcomes.

