

7. Daxboek F, Zitta S, Assadian O, Krause R, Wenish C. *Ochrobactrum anthropi* bloodstream infection complicating hemodialysis. *Am J Kidney Dis.* 2002;40:e17.1-4.
8. Chertow G. *Ochrobactrum anthropi* bacteremia in a patient on hemodialysis. *Am J Kidney Dis.* 2000;35:e30.1-2.
9. Soloaga R, Carrión N, Pidone J, Guelfand L, Margari A, Altieri R. Bacteriemia relacionada a catéter por *Ochrobactrum anthropi*. *Medicina (B Aires).* 2009;69:655-7.
10. Ospina S, Atehortúa Muñoz S, Zapata J. Bacteriemia por *Ochrobactrum anthropi* en paciente con obstrucción de la vía biliar. *Infect.* 2009;13:293-5.

Esther Torres Aguilera^{a,*}, Eduardo Verde Moreno^a,
Patricia Muñoz^b, Maricela Valerio^b, Jose Luño^a

^a Servicio de Nefrología, Hospital General Universitario Gregorio Marañón, Madrid, Spain

^b Servicio de Microbiología, Hospital General Universitario Gregorio Marañón, Madrid, Spain

* Corresponding author.

E-mail address: estertor.ag@gmail.com (E. Torres Aguilera).

<http://dx.doi.org/10.1016/j.nefro.2017.01.019>

2013-2514/© 2016 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Hypercalcemia and hypophosphatemia in a 11 years old girl with ovarian tumour[☆]

Hipercalcemia e hipofosforemia en niña con tumour ovárico

Dear Editor,

The most common cause of hypercalcaemia in patients with non-metastatic solid tumours is the secretion of parathyroid hormone-related (PTHr) protein.¹

The patient is an 11-year-old female with no history of interest, to be evaluated for an abdominal mass. Upon arrival the patient showed high serum creatinine, hypercalcaemia and hypophosphatemia (Table 1). The baseline renal function test also showed hypercalciuria and hyperphosphaturia. CT scan showed a large intra-abdominal mass. A complete resection of the tumour was

performed. An anatomopathological study revealed the presence of an ovarian granulosa cell tumour.

Clinical course was favourable. In the follow-up, one week after removal of the tumour, blood calcium levels normalised but urinary calcium remained high. The hypophosphataemia persisted, despite greater renal reabsorption of phosphate. After one month, all blood and urine biochemical parameters were within the normal range (Table 1).

Hypercalcaemia due to hypersecretion of PTHr by malignant tumours is an exceptional finding at paediatric age.^{2,3} Parathyroid hormone (PTH) and PTHr have shown similar effects in the renal handling of calcium and phosphate.⁴⁻⁷

Table 1 – Biochemical blood and urine values prior to tumour removal and at days 6, 30 and 60 after surgery.

	Pre-operative	Six days after surgery	Thirty days after surgery	Sixty days after surgery
Blood creatinine (mg/dl)	1.13	0.75	0.66	0.64
Blood-ionised calcium (mMol/l)	1.66	1.18	1.27	1.22
Blood phosphate (mg/dl)	2.6 mgr/dl	2.8 mgr/dl	5.2	5.4
PTH intact (pg/ml)	2.7	44	–	–
PTHr ^a (pmol/l)	10.8	<1.1	–	–
Vitamin D (ng/ml)	30	–	–	–
Calcitonin (pg/ml)	10.4	–	–	–
Urine Ca/Cr ratio	1.4	0.46	0.09	0.07
CEI ^a	1.59	0.34	0.06	0.05
Calciuria (mg/kg/24 h)	15	7.43	1.61	1.26
TRP ^a	32	96	94	90
TP/GFR ^a	0.85	2.71	4.88	4.82
Phosphaturia (mg/24 h/1.73 m ²)	954	110	427	880

CEI: calcium excretion index; PTHr: parathyroid hormone-related protein; TP/GFR: phosphate per 100 ml of glomerular filtration rate; TRP: tubular reabsorption of phosphate.

^a Source: Santos and García-Nieto.⁹

DOI of original article:

<http://dx.doi.org/10.1016/j.nefro.2016.09.008>.

[☆] Please cite this article as: Ubetagoyena Arrieta M, Martínez Sainz de Jubera J, García de Andoin Barandiaran N, Úriz Monaut JJ, López Cuesta S, Domínguez Castells A. Hipercalcemia e hipofosforemia en niña con tumour ovárico. *Nefrologia.* 2017;37:100-101.

They release calcium from bone and promote the renal reabsorption of calcium in the distal nephron. They also produce hyperphosphaturia by decreasing the tubular reabsorption of phosphate that may result in hypophosphataemia. However, patients with hyperparathyroidism may present with hypercalciuria.⁸ The elevated calcaemia in our patient is related to high levels of PTHr. Significantly, hypercalciuria persisted after excision of the tumour and normalisation of calcaemia levels.

Another remarkable finding was the presence of hypophosphataemia, which became significant. The hypophosphataemia coincided with a situation of renal phosphate loss, evidenced by low levels of TRP and TP/GFR, indicating that the hypophosphataemia was of renal origin.

In conclusion, we present a case of PTHr-mediated hypercalcaemia secondary to an ovarian tumour in an age group that is rarely affected by this condition. In addition to transient renal failure, the biochemical data show hypercalcaemia accompanied by hypercalciuria. The hypophosphataemia is secondary to an excessive loss of phosphate through the kidney.

REFERENCES

1. Nakamura Y, Bando Y, Shintani Y, Yokogoshi Y, Saito S. Serum parathyroid hormone-related protein concentrations in patients with hematologic malignancies or solid tumors. *Acta Endocrinol (Copenh)*. 1992;127:324-30.
2. Piura B, Wiznitzer A, Shaco-Levy R. Juvenile granulosa cell tumor of the ovary associated with hypercalcemia. *Arch Gynecol Obstet*. 2008;277:257-62.
3. Daubenton JD, Sinclair-Smith C. Severe hypercalcemia in association with a juvenile granulosa cell tumor of the ovary. *Med Pediatr Oncol*. 2000;34:301-3.
4. Horwitz MJ, Tudesco MB, Sereika SM, Hollis BW, Garcia-Ocaña A, Stewart AF. Direct comparison of sustained infusion of human parathyroid hormone-related protein-(1-36) [hPTHrP-(1-36)] versus hPTH-(1-34) on serum calcium, plasma 1,25-dihydroxyvitamin D concentrations, and fractional

calcium excretion in healthy human volunteers. *J Clin Endocrinol Metab*. 2003;88:1603-9.

5. Gesek FA, Friedman PA. On the mechanism of parathyroid hormone stimulation of calcium uptake by mouse distal convoluted tubule cells. *J Clin Invest*. 1992;90:749-58.
6. Syed MA, Horwitz MJ, Tedesco MB, Garcia-Ocaña A, Wisniewski SR, Stewart AF. Parathyroid hormone-related protein-(1-36) stimulates renal tubular calcium reabsorption in normal human volunteers: Implications for the pathogenesis of humoral hypercalcemia of malignancy. *J Clin Endocrinol Metab*. 2001;86:1525-31.
7. Esbrit P. Mecanismo de acción de la PTH en el túbulo renal. *Nefrología*. 1988;1:4-8.
8. Clines GA. Mechanisms and treatment of hypercalcemia of malignancy. *Curr Opin Endocrinol Diabetes Obes*. 2011;18:339-44.
9. Santos F, García-Nieto V. Función renal basal. In: García Nieto V, Santos F, editors. *Nefrología Pediátrica*. 2.ª edición Madrid: Grupo Aula Médica; 2006. p. 39-49.

Mercedes Ubetagoyena Arrieta^{a,*},
Jorge Martínez Sainz de Jubera^a,
Nagore García de Andoin Barandiaran^b,
José Javier Úriz Monaut^b, Sheila López Cuesta^a,
Ana Domínguez Castells^b

^a Sección de Nefrología Pediátrica, Hospital Universitario Donostia, San Sebastián, Guipúzcoa, Spain

^b Servicio de Oncología Pediátrica, Hospital Universitario Donostia, San Sebastián, Guipúzcoa, Spain

*Corresponding author.

E-mail address:

m.mercedes.ubetagoyenaarrieta@osakidetza.net

(M. Ubetagoyena Arrieta).

<http://dx.doi.org/10.1016/j.nefro.2017.01.023>

2013-2514/© 2016 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Diagnosis of obstruction peritoneal catheter by fluoroscopic peritoneography[☆]

Diagnóstico de obstrucción de catéter peritoneal mediante peritoneografía fluoroscópica

Dear Editor,

Infusion/drainage problems are defined as a slowed flow, hampered or prevented by causes related to the catheter itself and not to the functioning of the peritoneum as a dialysis

membrane. The occurrence of these problems ranges between 5% and 20% and is often associated with the technique used for the placement of the catheter. They are more common in cases of implantation by laparoscopy.¹

[☆] Please cite this article as: Sastre A, González-Arregoces J, Romainoik I, Mariño S, Lucas C, Monfá E, et al. Diagnóstico de obstrucción de catéter peritoneal mediante peritoneografía fluoroscópica. *Nefrología*. 2017;37:101-103.