



Original

Serum ICAM-1 level and ICAM-1 gene 1462A>G (K469E) polymorphism on microalbuminuria in nondiabetic, nonhypertensive and normolipidemic obese patients: Genetical background of microalbuminuria in obesity

Ahmet Engin Atay^a, Bennur Esen^{b,*}, Halit Akbas^c, Emel Saglam Gokmen^a, Saadet Pilten^d, Hale Guler^c, Dilek Gogas Yavuz^e

^a Division of Nephrology, Department of Internal Medicine, Bagcilar Research and Training Hospital, Istanbul, Turkey

^b Division of Nephrology, Department of Internal Medicine, Acibadem University School of Medicine, Istanbul, Turkey

^c Department of Medical Biology and Genetics, Medical School of Harran University, Sanliurfa, Turkey

^d Department of Biochemistry, Bagcilar Research and Training Hospital, Istanbul, Turkey

^e Department of Endocrinology and Metabolism, Marmara University School of Medicine, Istanbul, Turkey

ARTICLE INFO

Article history:

Received 14 April 2016

Accepted 17 November 2016

Available online 30 May 2017

Keywords:

Obesity

Microalbuminuria

Adhesion molecule

ICAM-1 gene

Polymorphism

ABSTRACT

Background: A growing body of evidence suggest that obese individuals are under risk of renal parenchymal disorders when compared to nonobese counterparts. Microalbuminuria is the early marker of renal involvement. Although most of obese patients carries multiple risk factors for microalbuminuria, some obese individuals without risk factor may progress to microalbuminuria. The present study was performed to examine the role of ICAM-1 gene 1462A>G (K469E) polymorphism on microalbuminuria in obese subjects without diabetes mellitus, hypertension, hiperlipidemia and older age.

Methods: Ninety eight obese and 96 nonobese individuals without a comorbidity enrolled into the study. Serum ICAM-1 level was measured by enzyme linked immunoabsorbent assay (ELISA) method. ICAM-1 gene 1462A>G (K469E) polymorphism was examined by restriction fragment length polymorphism-polymerase chain reaction (RFLP-PCR). Nephelometric method was used to examine urinary albumin loss, and microalbuminuria was measured by albumin to creatinine ratio.

Results: Obese individuals had significantly higher microalbuminuria and proteinuria level compared to nonobese subjects (p : 0.043 and p : 0.011; respectively). GG genotype of ICAM-1 carriers have significantly higher microalbuminuria compared to individuals with AA or AG genotype carriers (p : 0.042). Serum ICAM-1 level was significantly correlated with creatinine

* Corresponding author.

E-mail address: bennuresen@yahoo.com (B. Esen).

<http://dx.doi.org/10.1016/j.nefro.2017.06.005>

2013-2514/© 2017 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

and microalbuminuria (p : 0.002 and p : 0.03; respectively). Logistic regression analysis indicated a 7.39 fold increased risk of microalbuminuria in individuals with GG genotype of ICAM-1 gene 1462A>G (K469E) polymorphism.

Conclusions: GG genotype of ICAM-1 gene K469E polymorphism is associated with increased microalbuminuria in obese individuals without another metabolic risk factor.

© 2017 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Polimorfismo 1462A>G (K469E) del gen ICAM-1 y nivel sérico de ICAM-1 en la oligoalbuminuria de pacientes obesos no diabéticos, no hipertensos y normolipidémicos: acervo genético de la oligoalbuminuria en la obesidad

R E S U M E N

Palabras clave:

Obesidad
Oligoalbuminuria
Molécula de adhesión
Gen ICAM-1
Polimorfismo

Introducción: Un conjunto de datos en aumento indica que los individuos obesos corren más riesgo de sufrir trastornos del parénquima renal si se los compara con sus homólogos no obesos. La oligoalbuminuria es un primer rasgo de afectación renal. Aunque la mayoría de los pacientes obesos presentan múltiples factores de riesgo de oligoalbuminuria, esta puede manifestarse en algunos individuos obesos sin factores de riesgo. El presente estudio se realizó para analizar el papel del polimorfismo 1462A>G (K469E) del gen ICAM-1 en la oligoalbuminuria de individuos obesos sin diabetes mellitus, hipertensión, hiperlipidemia ni vejez.

Métodos: Para el estudio fueron reclutados 98 individuos obesos y 96 individuos no obesos sin comorbilidad. Se midió el nivel sérico de ICAM-1 mediante el ensayo de inmunoabsorción enzimática (ELISA). Se analizó el polimorfismo 1462A>G (K469E) del gen ICAM-1 por reacción en cadena de la polimerasa y polimorfismo de longitud de los fragmentos de restricción (RFLP-PCR). El método nefolométrico se utilizó para analizar la pérdida urinaria de albúmina, y la oligoalbuminuria se midió con la tasa de albúmina/creatinina.

Resultados: Los individuos obesos presentaron unos niveles de oligoalbuminuria y proteinuria considerablemente más elevados que los individuos no obesos (p : 0,043 y p : 0,011, respectivamente). La oligoalbuminuria en los portadores del genotipo GG de ICAM-1 fue bastante mayor que la de los portadores del genotipo AA o AG (p : 0,042). El nivel sérico de ICAM-1 se correlacionó notablemente con la creatinina y la oligoalbuminuria (p : 0,002 y p : 0,03, respectivamente). El análisis de regresión logística mostró un riesgo 7,39 veces mayor de oligoalbuminuria en los individuos con el genotipo GG del polimorfismo 1462A>G (K469E) del gen ICAM-1.

Conclusiones: El genotipo GG del polimorfismo 1462A>G (K469E) del gen ICAM-1 se asocia con un aumento de la oligoalbuminuria en personas obesas sin otro factor de riesgo metabólico.

© 2017 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Obesity is becoming worldwide epidemic which accounts a risk factor for metabolic and hormonal disorders.¹ Because insulin resistance is frequently seen, obese individuals are considered as candidates of diabetes mellitus (DM) and associated microvascular complications.² Additionally, increased adipose tissue is also a source of subclinic inflammation which is closely related to endothelial damage.

Microalbuminuria (MA) is initial stage of diabetic nephropathy which is strongly correlated with endothelial injury.³ MA comprise an independent risk factor for coronary artery disease in both diabetic and nondiabetic population.⁴ Obese individuals even without overt DM and hypertension may exhibit microalbuminuria.^{5,6} Enhanced blood flow and

glomerular filtration rate (GFR) are resulted with MA. An increasing number of evidence indicates low-grade inflammation as a cause of MA.⁷ Also obesity is a well-known risk factor for focal segmental glomerulosclerosis.⁸

One of the suggested mechanisms in the pathogenesis of microalbuminuria is overexpression of adhesion molecules including ICAM-1 which closely associated with severe inflammatory processes that usually resulted with tissue injury.⁹ Patients with obesity and diabetic nephropathy exhibit overexpression of ICAM-1.^{10,11} Experimental studies showed that hyperglycemia is associated with overexpression of ICAM-1 in renal glomerular tissue and renal endothelial injury.¹² However, varying degree of renal injury among patients with normal ICAM-1 level or disassociation of ICAM-1 level with microalbuminuria may suggest the role of genetic mutations in ICAM-1 gene. As a marker of cell integrity and complexity,

the effect of ICAM-1 on renal tissue is well-documented however lack of available data exist regarding to the role of ICAM-1 gene 1462A>G (K469E polymorphism on microalbuminuria in obese individuals.

Our aim is to examine the role of serum ICAM-1 level and ICAM-1 gene 1462A>G polymorphism on urine albumin loss in metabolically normal obese and nonobese individuals.

Material and methods

A total of 194 participants; 98 obese and 96 nonobese, older than 18 years without a history of systemic disorder that admitted to obesity outpatient service of Bagcilar Education and Research Hospital between February 2015 and June 2015 were enrolled. Height and weight of participants were measured in Tanita Body Composition Analyzer (Tanita Corporation of America, Illinois, USA). Body mass index (BMI) was calculated as weight/height² (kg/m²). Body mass index (BMI) ≥ 30 kg/m² was defined as obesity.¹³ Exclusion criterias were secondary obesity, fasting glucose ≥ 100 mg/dl, HbA1c ≥ 5.8 , hypertension, hyperlipidemia, CVD, renal parenchymal disorder or urolitiazis.¹⁴ Hypertension was defined as systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg, or receiving any antihypertensive agent. Renal parenchymal disorder or urolitiazis was diagnosed by ultrasonographic examination. National Cholesterol Education Programme (NCEP) was used to define hyperlipidemia (LDL-cholesterol > 130 mg/dl or trygliceride > 200 mg/dl).¹⁵ Ethics committee of Bagcilar Education and Research Hospital approved the study. Written informed consent was obtained from all participants. The study was performed in accordance with Declaration of Helsinki. Clinical Trials Registration number was 09.2014.0310. Registration date was 16/01/2015

Sample collection and analysis

Blood samples were obtained after 8-h fasting period. Serum ICAM-1 concentration was analyzed by enzyme linked immunosorbent assay (ELISA) by commercially available kits (Invitrogen, Thermo Fisher Scientific, USA). Whole blood count was performed in blood samples that were taken into tubes with EDTA using an automatic blood counter (XE-5000; Sysmex Corp, Kobe, Japan). Biochemical variables including serum glucose, urea, creatinine, uric acid, aspartat aminotransferase (AST), alanine aminotransferase (ALT), lactate dehydrogenase (LDH), calcium, sodium, potassium, clor, total protein, albumin, total cholesterol, low density lipoprotein (LDL), high density lipoprotein (HDL), trigliceride, were analyzed by photometric method in Siemens Advia 1800 device (Siemens Healthcare Diagnostics, Kobe, Japan). Hormon parameters including HbA1c, insulin, C-peptide, TSH were analyzed by chemiluminescence immunoassay method in Siemens Advia Centaur device (XE-5000, Sysmex Corp., Kobe, Japan).

Urine analysis was performed by spectrophotometric method in Siemens Advia 1800 device (XE-5000, Sysmex Corp., Kobe, Japan). Urinary albumin was measured by immunoturbidimetric method in Cobas auto analyzer (Roche Diagnostics, Germany). Urinary creatinine was analyzed by Aeroset auto-analyzer (Abbott Laboratories Inc., Abbott Park, IL, USA). Urine

albumin excretion rate exceeding 20 μ g/min (30 mg/day); in the absence of uncontrolled hypertension or urinary tract infection, was defined as microalbuminuria.

DNA extraction

Total genomic DNA was extracted from peripheral blood leucocytes using a purification kit (Thermo Scientific GeneJET™ Whole Blood Genomic DNA Purification Kit) according to previously described.¹⁶

Genotyping analysis

To analyze ICAM-1 gene 1462A>G polymorphism (K469E; rs5498 polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method was. PCR amplification was generated using the following oligonucleotide primers: 5'-GGAACCCATTGCCCGAGC-3' and reverse 5'-GGTGAGGATTGCATTAGGTC-3' (product of 223 bp).

A PCR reaction was carried out in a 10 μ l reaction volume containing 1 \times PCR buffer, 2 mM MgCl₂, 0.2 mM each deoxynucleotide triphosphate (dNTPs, Fermentas, St. Leon-Rot, Germany), 40 ng of DNA, 0.2 μ M of each primer (Bio Basic Inc., Ontario, Canada), and 0.5 unit of Taq DNA Polymerase (Fermentas). The PCR conditions were: 3 min of initial denaturation at 94 °C, followed by 30 cycles at 95 °C for 30 s, 30 s at 65 °C for annealing, and 30 s at 72 °C for extension, followed by 5 min at 72 °C for final extension. The PCR products were 223 bp.

For the RFLP analysis, PCR-amplified products were digested with BanII for 2–5 h at 37 °C (New England Biolabs Inc., Hertfordshire, UK). The amplicon with the homozygous GG allele of ICAM was cleaved by BanII, yielding 136 and 87 bp fragments, whereas the amplicon with the homozygous AA allele remained uncut, yielding a 223 bp band. The amplicon with the heterozygous AG allele yielded three fragments of 223, 136 and 87 bp length. The digested products were separated on 3% agarose gel along with a 100–1500 bp DNA ladder (Bio Basic Inc.). Ethidium bromide-stained gels were visualized under UV light using the Alpha Imager System (Alpha Innotech, San Leandro, CA, USA) (Fig. 1).

Statistical analysis

Data were analyzed using the SPSS for Windows computer program (release 22.0; SPSS Inc. Chicago, IL, USA). All data




Fig. 1 – 2% agarose gel electrophoresis stained with ethidium bromide showing the PCR-RFLP analysis of the ICAM-1. Lanes M: DNA marker. Lanes 1, 2: AA allelic polymorphism; Lane 3, 4: AG allelic polymorphism; Lane 5, 6: GG allelic polymorphism.

Table 1 – Comparison of demographic characteristics and laboratory data between obese and nonobese individuals.

	Obese (n = 98)	Nonobese (n = 96)	p
	Mean ± SD	Mean ± SD	
Age (years)	37.8 ± 9.79	37.77 ± 8.83	NS
Glucose	93.4 ± 6.11	92.98 ± 7.87	NS
Urea	27.4 ± 8.1	27.11 ± 8.24	NS
Creatinin	0.94 ± 1.57	0.74 ± 0.15	NS
Uric acid	5.29 ± 1.5	4.27 ± 1.04	^a 0.001**
AST	23.35 ± 14.44	17.94 ± 4.36	^b 0.038*
ALT	33.31 ± 38.83	16.64 ± 8.56	NS
LDH	204.28 ± 32.6	187.85 ± 32.17	^a 0.001**
Trygliceride	146.54 ± 90.32	114.54 ± 73.55	NS
Cholesterol	187.02 ± 41.79	182.47 ± 32.32	NS
HDL-c	52.01 ± 42.51	54.91 ± 14.08	NS
LDL-c	111.14 ± 34.83	105.51 ± 30.22	NS
Ca	9.54 ± 0.48	9.45 ± 0.79	NS
Na	140.51 ± 14.22	141.45 ± 2.26	NS
K	4.52 ± 0.28	4.4 ± 0.36	^a 0.008**
Cl	113 ± 9.3	104 ± 10.77	^b 0.027*
CRP	5.07 ± 4.44	2.55 ± 5.37	NS
T. protein	7.4 ± 0.41	7.39 ± 0.44	NS
Albumin	4.56 ± 0.31	4.66 ± 0.36	^a 0.039*
TSH	2.37 ± 2.37	2 ± 2.24	NS
S T4	1.13 ± 0.15	1.21 ± 0.18	^a 0.001**
Insulin	22.66 ± 26.24	9.74 ± 6.35	NS
C peptide	3.68 ± 1.85	2.37 ± 0.96	NS
Leukocyte count	8.03 ± 1.98	8.4 ± 1.73	NS
Hemoglobine	13.76 ± 1.89	13.72 ± 1.67	NS
Platelet count	282.01 ± 69.74	254.76 ± 59.28	^a 0.004**
Mean platelet volume	7.74 ± 1.22	8.24 ± 1.82	NS
Urine density	1018.59 ± 5.27	1006.07 ± 10.39	NS
Urine protein	0.15 ± 0.36	0.04 ± 0.19	^b 0.011*
Spot creatinin	186.5 ± 83.71	165.9 ± 83.24	NS
Spot MA/Cr	9 ± 13.17	6.58 ± 7.08	NS
Spot microalbumin	1.94 ± 4.31	1.16 ± 1.65	^b 0.043*
Serum ICAM-1	293.1 ± 231.4	250.7 ± 152.2	^a 0.001**

^a Student t test.
^b Mann Whitney U test.
* p < 0.05.
** p < 0.01.
Significance of bold values are provided.

Table 2 – Clinical and laboratory characteristics of obese group according to genotypes.

	ICAM-1			p
	AA (n = 26)	AG (n = 48)	GG (n = 24)	
	Mean ± SD	Mean ± SD	Mean ± SD	
Age (years)	36.97 ± 9.83	37.18 ± 8.74	37.92 ± 9.12	NS
BMI	28.95 ± 7.23	29.26 ± 6.45	29.87 ± 8.13	NS
Systolic blood pressure	122.87 ± 13.32	123.46 ± 13.81	125.78 ± 13.21	NS
Diastolic blood pressure	74.63 ± 9.87	77.08 ± 10.22	77.13 ± 8.89	NS
Glucose	95.12 ± 11.36	93.39 ± 10.29	95.32 ± 12.02	NS
Urea	28.16 ± 9.47	26.32 ± 6.71	28.43 ± 7.18	NS
Creatinin	0.71 ± 0.12	0.89 ± 0.37	0.78 ± 0.19	NS
Uric acid	4.56 ± 1.17	4.71 ± 1.38	5.37 ± 1.71	^a 0.038*
T.protein	7.43 ± 0.45	7.38 ± 0.41	7.41 ± 0.43	NS
Albumin	4.61 ± 0.41	4.57 ± 0.29	4.63 ± 0.24	NS
Urine density	1015.56 ± 5.91	1007.35 ± 6.02	1018.76 ± 5.42	NS
Urine protein	0.08 ± 0.26	0.04 ± 0.18	0.19 ± 0.39	^b 0.014*
Spot creatinin	184.78 ± 86.4	171.36 ± 78.97	190.67 ± 87.86	NS
Spot MA/Cr	7.76 ± 14.33	7.68 ± 9.23	8.68 ± 8.78	NS
Spot microalbumin	1.61 ± 1.58	1.34 ± 1.84	1.93 ± 2.14	^b 0.042*

^a Student t test.
^b Mann Whitney U test.
* p < 0.05.
** p < 0.01.

expressed as the mean \pm SD. Differences between the study groups were analyzed by using Student's t-test. Quantitative data were analyzed by chi-square test. Pearson correlation analysis and Spearman's rho correlation analysis were used to evaluate the correlation of variables. The variables significant in univariate analysis were included to multivariate logistic regression analysis (only significant correlation coefficients are reported). A *p* value <0.05 was considered to be statistically significant.

Findings

Table 1 shows demographic characteristics and laboratory parameters of obese and nonobese participants. Mean age of groups were similar however obese individuals had significantly higher uric acid, platelet count, microalbuminuria and proteinuria level (*p*: 0.001, *p*: 0.004, *p*: 0.043 and *p*: 0.011; respectively) (Table 1). Serum ICAM-1 level of obese individuals was significantly higher (*p*: 0.001).

Table 2 demonstrates comparison of different genotypes of ICAM-1 gene with regard to demographic characteristics and laboratory parameters. Obese individuals with GG genotype had significantly higher uric acid, microalbuminuria and proteinuria when compared to individuals with AA or AG genotype (*p*: 0.038, *p*: 0.042 and *p*: 0.014; respectively) (Table 2).

Male participants had significantly higher urea, creatinine, AST, ALT, LDH, triglyceride, total protein, albumin, FT4, hemoglobin and urine creatinine level while female participants had higher levels of body fat ratio and platelet count (*p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.023, *p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.001 and *p*: 0.002; respectively).

Correlation analysis indicated a significant correlation between serum ICAM-1 with HbA1c, creatinine, uric acid, AST, ALT, LDH, total cholesterol, LDL-c, albumin, hemoglobin and microalbuminuria (*p*: 0.027, *p*: 0.002, *p*: 0.001, *p*: 0.011, *p*: 0.001, *p*: 0.006, *p*: 0.043, *p*: 0.012, *p*: 0.046, *p*: 0.001 and *p*: 0.030; respectively) (Table 3). Association of creatinine with urea, uric acid, AST, ALT, LDH, albumin, TSH, FT4, WBC, hemoglobin, platelet count, urine creatinine and body fat mass were significant (*p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.001, *p*: 0.024, *p*: 0.004, *p*: 0.001, *p*: 0.001, *p*: 0.001 and *p*: 0.001; respectively).

Multivariate analysis and Logistic regression analysis were performed to predict microalbuminuria risk in obese individuals with different genotypes of ICAM-1 gene single nucleotide polymorphism and determined that obese individuals with GG genotype of ICAM-1 gene 1462A>G polymorphism have 7.39 fold increased risk of microalbuminuria (*p*: 0.049) while the risk of microalbuminuria in individuals with AA or AG genotype were nonsignificant (Tables 4 and 5).

Discussion

The present study indicated that obese individuals have worse metabolic control and evident microalbuminuria when compared to nonobese counterparts. Female individuals had significantly higher body fat ratio however metabolic deterioration of male individuals was evident. GG genotype

Table 3 – Correlation of serum ICAM-1 with demographic and laboratory variables.

	ICAM-1	
	<i>r</i>	<i>p</i>
Age (years)	-0.137	NS
BMI	0.096	NS
Metabolic age (years)	-0.085	NS
Systolic blood pressure	0.003	NS
Diastolic blood pressure	-0.078	NS
HbA1c	0.162	0.027*
Glucose	-0.010	NS
Urea	0.080	NS
Creatinin	0.222	0.002**
Uric acid	0.297	0.001**
AST	0.183	0.011*
ALT	0.264	0.001**
LDH	0.197	0.006**
Trygliceride	0.106	NS
Cholesterol	0.148	0.043*
HDL-c	-0.114	NS
LDL-c	0.182	0.012*
Ca	0.198	0.006**
Na	0.127	NS
K	0.007	NS
GI	-0.043	NS
CRP	0.092	NS
T.protein	0.165	0.023*
Albumin	0.145	0.046*
TSH	0.083	NS
S T4	0.112	NS
Insulin	0.059	NS
C peptide	0.067	NS
Leukocyte count	0.090	NS
Hemoglobine	0.243	0.001**
Hematocryte	0.230	0.001**
Platelet count	-0.015	NS
Mean platelet volume	-0.146	0.044*
Urinedensity	-0.059	NS
Urine protein	-0.051	NS
Spot creatinin	-0.008	NS
Spot MA/Cr	-0.030	NS
Spot microalbumin	-0.035	0.030*
Spearman Rho test.		
* <i>p</i> < 0.05.		
** <i>p</i> < 0.01.		

of ICAM-1 gene is associated with microalbuminuria and proteinuria. Finally, we determined a significant correlation between serum ICAM-1 level with microalbuminuria and metabolic control.

Endothelial dysfunction is the initial step of vascular changes which eventually progress to evident atherosclerosis (AS) and AS related complications.¹⁷ Endothelial injury and related disorders are five times more frequent and significantly seen at earlier ages among obese population.¹ For more than a century ago, relation of obesity and AS has been known.¹⁸ Obesity is a well-known criteria of metabolic syndrome (MS) and accounts four times higher risk for atherosclerotic disorders.¹⁹ In a study including 9961 obese subjects, it has been shown that obesity is risk factor for microalbuminuria independent from diabetes and hypertension.²⁰ In agreement with the literature, our obese population had

Table 4 – Multivariate analysis of different ICAM-1 genotypes.

	B	S.E.	p	Exp(B)	95.0% C.I. for Exp(B)	
					Lower	Upper
ICAMgr2			0.048			
ICAM AG	-1.08	0.48	0.042	0.341	0.133	0.875
ICAM GG	-0.84	0.44	0.025	0.432	0.179	1.042

Variable(s) entered on step 1: ICAMgr2.

Table 5 – Logistic regression analysis to predict microalbuminuria in different genotypes of ICAM-1 gene K469E polymorphism.

	<0.3 Microalbuminuria		>0.3 Microalbuminuria		p	OR %95 GA
Obesity group						
AA	7	31.82%	18	25.35%		
AG heterozygote	14	63.64%	34	47.89%	0.91	0.94 (0.32–2.76)
GG homozygote	1	4.55%	19	26.76%	0.049	7.39 (0.83–66.2)
AG + GG	15	68.18%	53	74.65%	0.58	1.37 (0.48–3.91)

higher blood pressure, lipid and uric acid levels, and excrete significantly higher amount of microalbumin and protein.

A growing body of evidence indicate that subclinical inflammation is evident in obese individuals which is closely related to AS and insulin resistance (IR).²¹ Inflammatory conditions are associated with increased endothelial activation and overexpression of adhesion molecules.²² Adhesion molecules regulates leukocyte migration across the endothelium and involved in integral part of inflammation.¹⁰ Obese individuals exhibit increased mRNA expression of ICAM-1 leading to infiltration of adipose tissue and other systems including kidneys by macrophages.²³ Independent from fat distribution, total fat volume is major determinant of serum levels of adhesion molecules.²⁴

Obese individuals or diabetic patients without well-controlled glycemia have increased ICAM-1 levels when compared to nonobese or diabetic patients with glycemic control.^{25,26} A well-known correlation has been established between diabetic complications including nephropathy and adhesion molecules.²⁷ A number of studies determined that patients with ICAM-1 gene K469E polymorphism, the K allele are significantly more prone to endothelial injury and CVD, and may predict future cardiovascular events.^{28–30} There are controversial reports regarding to relation of ICAM-1 gene polymorphism and microvascular complications of DM. Sun et al. showed that ICAM-1 gene polymorphism has nonsignificant effect on diabetic retinopathy (DR) while Liu et al. determined an increased risk of DR in patients with ICAM-1 gene polymorphism.^{31,32}

Increased serum levels of adhesion molecules may considered as early markers of endothelial injury which is resulted with systemic vascular disorder including renal microvasculature.^{33,34} There is a positive correlation between serum ICAM-1 level and ICAM-1 gene polymorphism.^{35,36} It would be reasonable to expect microalbuminuria in obese individuals with overexpression of adhesion molecules because obese individuals have impaired insulin sensitivity, and also adhesion molecules destruct beta-cells leading to insulinitis, hyperglycemia and hyperglycemia related nephrotoxicity that all may resulted with microalbuminuria.

Adhesion molecules including ICAM-1 have been demonstrated in atherosclerotic lesions which is regressed by adiponectin, a adipocyte secretory protein decreased in obesity.³⁷

The present study has some potential drawbacks. First, sample size of the study was relatively low. We included relatively young obese and nonobese individuals without diabetes, hypertension, hyperlipidemia, renal parenchymal or cardiovascular disorders to eliminate cumulative effect of these entities on microalbuminuria. Second, this study was conducted in Turkish population that limits us to generalize our results to other populations. Finally, evaluation of complex interaction of weight loss and serum ICAM-1 level may provide complementary data on microalbuminuria in obesity.

The importance/novelty of the study is that it is the first report that examined the role of ICAM-1 gene 1462A>G polymorphism on microalbuminuria in obesity, and established that GG genotype of ICAM-1 gene is an independent risk factor of microalbuminuria in obesity. There is a limited data on the genetic background of microalbuminuria in obese individuals. Understanding molecular and genetic mechanisms of microalbuminuria may help to identify obese patients under risk of nephropathy that is initially manifested with microalbuminuria.

Based on the fact that ICAM-1 is involved in endothelial injury and atherosclerosis, synthesis and genetic basis of this molecule may be therapeutic target for the complications of obesity. Further studies with large sample size are warranted to reach more precise conclusion.

Ethical disclosure

The study was performed in accordance with Helsinki Declaration.

Ethics Committee approval was obtained.

Informed consent was obtained from all participants.

Conflict of interests

Authors declare that there is no conflict interest.

REFERENCES

1. Bray AG. Health hazard of obesity. *Endocrinol Metab Clin North Am.* 1996;25:907-18.
2. WHO, Obesity and overweight Fact sheet No. 311; May 2012. URL: <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>
3. Pontremoli R, Nicoletta C, Viazzi F, Ravera M, Sofia A, Berruti V, et al. Microalbuminuria is an early marker of target organ damage in essential hypertension. *Am J Hypertens.* 1998;11:430-8.
4. Ekicibas E, Kaderli AA, Kazazoglu AR, Ozdemir B. Relation of coronary artery disease severity with microalbuminuria and fasting insulin level in nondiabetic individuals. *Anatolian Cardiol J.* 2008;8:16-21.
5. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification; 2012. URL: http://www.kidney.org/professionals/kdoqi/guidelines_ckd/toc.htm
6. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976;16:31-41.
7. De Jong PE, Verhave JC, Pinto-Sietsma SJ, Hillege HL, PREVEND study group. Obesity and target organ damage: kidney. *Int J Obes Relat Metab Disord.* 2002;26:21-4.
8. Kambham N, Markowitz GS, Valeri AM, Lin J, D'Agati VD. Obesity-related glomerulopathy: an emerging epidemic. *Kidney Int.* 2001;59, 1498-1509.
9. Mitsui A, Tada Y, Shibata S, Kamata M, Hau C, Asahina A, et al. Deficiency of both L-selectin and ICAM-1 exacerbates imiquimod-induced psoriasis-like skin inflammation through increased infiltration of antigen presenting cells. *Clin Immunol.* 2015, pii:S1521-6616(14)00289-7.
10. Jamar G, Pisani LP, Medeiros A, Oyama LM, Masquiao DC, Colantonio E, et al. Effect of fat intake on the inflammatory process and cardiometabolic risk in obesity after interdisciplinary therapy. *Horm Metab Res.* 2016;48:106-11.
11. Mudaliar H, Pollock C, Ma J, Wu H, Chadban S, Panchapakesan U. The role of TLR2 and 4-mediated inflammatory pathways in endothelial cells exposed to high glucose. *PLoS One.* 2014;9:e108844.
12. Soares BL, de Freitas MA, Montero EF, Pitta GB, Miranda F Jr. Alprostadil attenuates inflammatory aspects and leucocytes adhesion on renal ischemia and reperfusion injury in rats. *Acta Cir Bras.* 2014;29 Suppl. 2:55-60.
13. WHO Expert Consultation. Appropriate body mass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004;363:157-63.
14. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2010;33 Suppl. 1:S62-9.
15. Expert Panel on Detection Evaluation and Treatment of High blood cholesterol in adults: third Report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III) final report. *Circulation.* 2002;106:3143-421.
16. Atay AE, Akbas H, Tumer C, Sakar MN, Incebiyik A, Esen B, et al. The association of endothelial nitric oxide synthase gene G894T polymorphism and serum nitric oxide concentration with microalbuminuria in patients with gestational diabetes. *Clin Nephrol.* 2013;81:105-11, <http://dx.doi.org/10.5414/CN108138>.
17. Allan RB, Vun SV, Spark JI. A comparison of measures of endothelial function in patients with peripheral arterial disease and age and gender matched controls. *Int J Vasc Med.* 2016;2016:2969740.
18. Curtin RG. The heart and its danger in the treatment of obesity. *Trans Am Climatol Assoc.* 1908;24:163-76.
19. Lee YA, Kang SG, Song SW, Rho JS, Kim EK. Association between metabolic syndrome, smoking status and coronary artery calcification. *PLoS One.* 2015;10:e0122430.
20. Rim JH, Lee YH, Cha BS, Lee SG, Kim JH. Central obesity is an independent risk factor for microalbuminuria in both the general Korean women and nondiabetic nonhypertensive subpopulation: association of microalbuminuria and metabolic syndrome from the Korea National Health and Nutrition Examination Survey 2011-2012. *Clin Chim Acta.* 2015;448:74-9.
21. Bosanská L, Michalský D, Lacinová Z, Dostálová I, Bártlová M, Haluzíková D, et al. The influence of obesity and different fat depots on adipose tissue gene expression and protein levels of cell adhesion molecules. *Physiol Res.* 2010;59:79-88.
22. Ara J, Mirapeix E, Arrizabalaga P, Rodriguez R, Ascaso C, Abellana R, et al. Circulating soluble adhesion molecules in ANCA-associated vasculitis. *Nephrol Dial Transplant.* 2001;16:276-85.
23. Hatunic M, Finucane F, Burns N, Gasparro D, Nolan JJ. Vascular inflammatory markers in early-onset obese and type 2 diabetes subjects before and after three months' aerobic exercise training. *Diab Vasc Dis Res.* 2007;4:231-4.
24. Brake DK, Smith EO, Mersmann H, Smith CW, Robker RL. ICAM-1 expression in adipose tissue: effects of diet-induced obesity in mice. *Am J Physiol Cell Physiol.* 2006;291: C1232-9.
25. Matsumoto K, Sera Y, Abe Y, Tominaga T, Horikami K, Hirao K, et al. High serum concentrations of soluble E-selectin correlate with obesity but not fat distribution in patients with type 2 diabetes mellitus. *Metabolism.* 2002;51:932-4.
26. Matsumoto K, Sera Y, Nakamura H, Ueki Y, Miyake S. Serum concentrations of soluble adhesion molecules are related to degree of hyperglycemia and insulin resistance in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract.* 2002;55:131-8.
27. Polat SB, Ugurlu N, Aslan N, Cuhaci N, Ersoy R, Cakir B. Evaluation of biochemical and clinical markers of endothelial dysfunction and their correlation with urinary albumin excretion in patients with type 1 diabetes mellitus. *Arch Endocrinol Metab.* 2016, pii:S2359-39972016005001104.
28. Liu ZP, Huo Y, Li JP, Zhang Y, Xue L, Zhao CY, et al. Polymorphism K469E of intercellular adhesion molecule-1 gene and restenosis after coronary stenting in Chinese patients. *Chin Med J (Engl).* 2004;117:172-5.
29. Zhang SR, Xu LX, Gao QQ, Zhang HQ, Xu BS, Lin J, et al. The correlation between ICAM-1 gene K469E polymorphism and coronary heart disease. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi.* 2006;23:205-7.
30. Lu FH, Shang Q, Wen PE, Su GH, Wu JM, Tian Q, et al. A study on K469E polymorphism of ICAM1 gene and ICAM1 plasma level in patients with coronary heart disease. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi.* 2006;23:195-7.
31. Sun H, Cong X, Sun R, Wang C, Wang X, Liu Y. Association between the ICAM-1 K469E polymorphism and diabetic retinopathy in Type 2 diabetes mellitus: a meta-analysis. *Diabetes Res Clin Pract.* 2014;104:e46-9.
32. Liu L1, Yu Q, Wang H, Zhang SX, Huang C, Chen X. Association of intercellular adhesion molecule 1 polymorphisms with retinopathy in Chinese patients with Type 2 diabetes. *Diabet Med.* 2006;23:643-8.
33. Glowinska B, Urban M, Peczyńska J, Florys B. Soluble adhesion molecules (sICAM-1, sVCAM-1) and selectins (sE selectin, sP selectin, sL selectin) levels in children and adolescents with obesity, hypertension, and diabetes. *Metabolism.* 2005;54:1020-6.
34. Głowińska-Olszewska B, Urban M, Tołwińska J, Peczyńska J, Florys B. Correlation analysis between biochemical and biophysical markers of endothelium damage in children with

- diabetes type 1. *Endokrynol Diabetol Chor Przemiany Materii Wieku Rozw.* 2005;11:221-7.
35. Singh A, Singh AK, Singh SK, Paliwal VK, Gupta RK, Prasad KN. Association of ICAM-1 K469E polymorphism with neurocysticercosis. *J Neuroimmunol.* 2014;276:166-71.
36. Wang MY, Bai DC, Zhu P, Fu Y, Bu DF, Zhang Y. Effects of ICAM-1 gene K469E, K56M polymorphisms on plasma sICAM-1 expression levels in Chinese Yugur, Tibetan and Han nationalities]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi.* 2012;20:1205-11.
37. Arita Y, Maeda K, Kuriyama H, Okamoto Y, Hotta K, Nishida M, et al. Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation.* 1999;100:2473-6.