

prednisone in steroid-resistant idiopathic membranoproliferative glomerulonephritis. *Clin Nephrol.* 2010;73:354-9.

6. Lim WH, Shingde M, Wong G. Recurrent and de novo glomerulonephritis after kidney transplantation. *Front Immunol.* 2019;10:1944.
7. Allen PJ, Chadban SJ, Craig JC, Lim WH, Allen RDM, Clayton PA, et al. Recurrent glomerulonephritis after kidney transplantation: risk factors and allograft outcomes. *Kidney Int.* 2017;92:461-9, <http://dx.doi.org/10.1016/j.kint.2017.03.015>.
8. Von Visger J, Cassol C, Nori U, Franco-Ahumada G, Nadasdy T, Satoskar AA. Complete biopsy-proven resolution of deposits in recurrent proliferative glomerulonephritis with monoclonal IgG deposits (PGNMIGD) following rituximab treatment in renal allograft. *BCM Nephrol.* 2019;20:53.

Leonidas Cruzado Vega*, Alba Santos García

Hospital General Universitario de Elche, Alicante, Spain

*Corresponding author.

E-mail address: leocruzadov@hotmail.com (L. Cruzado Vega).

2013-2514/© 2021 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
<https://doi.org/10.1016/j.nefro.2021.11.003>

Clinical utility of hybrid (SPECT/CT) imaging in calciphylaxis

Utilidad clínica de las imágenes híbridas (SPECT/CT) en la calcifilaxis

Dear Editor,

We congratulate Herrera-Martínez et al. for the excellent management of their patient with calciphylaxis.¹ With regards to the role of hybrid imaging in calciphylaxis we would like to mention few additional points which we believe could add to the fund of knowledge of our readers.

The important question to the research bodies working on calciphylaxis will be would hybrid image (SPECT/CT) replace skin biopsy. As suggested by Herrera-Martínez et al., skin biopsy has its own limitations and complications and any test like SPECT/CT hold that potential. However, we want to bring into the discussion regarding the close differentials of calciphylaxis. Will SPECT/CT be able to differentiate the various skin lesions like cellulitis, vasculitis, or atherosclerosis or warfarin necrosis? Other questions that require immediate attention are: cost effectiveness and clinical utility of early vs late SPECT/CT, clinical benefit of combined diagnostic approach (SPECT/CT plus skin biopsy) versus SPECT/CT alone, random skin biopsy of skin lesions vs SPECT/CT guided biopsy of the most intense uptake site. Most common site of calciphylaxis is lower limbs which is usually amenable site for skin biopsy. There are specific locations like penis where skin biopsy is contraindicated.^{2,3} In such locations, we believe that the clinical utility of SPECT/CT alone could potentially of extreme importance. Successful treatment by Herrera-Martínez et al. also encourages us to think that is there any correlation between the differential responses to therapy depending on extent of calciphylactic lesions detected on SPECT/CT?

Due to lower specificity of SPECT/CT, should we limit this study only to long-term non-healing ulcers in the setting of

end-stage renal failure? We highlight this point due to the fact that there are many other clinical differentials for the skin lesions in ESRD patients and doing a SPECT/CT in all cases may not be justified. The skin lesions in Herrera-Martínez et al. case was just few days old for which SPECT/CT was done. We wish to educate our readers not to make any general inference to do SPECT/CT in all cases of newly evolved skin lesions in ESRD. Ideal would be to diligently rule out all the other differentials as mentioned above.⁴ Another point on which future studies can be directed is to study the follow up SPECT/CT to see the follow up response of calciphylaxis to treatment of sodium thiosulphate.

In conclusion, newer non-invasive advancements in detecting calciphylaxis are encouraging but there are still many questions that needs to be settled before any guidelines can be laid down.⁵

Ethical statement

The article does not contain the participation of any human being and animal.

Funding

None declared.

Conflict of interest

Authors have no conflicts of interest to declare.

REFERENCES

1. Herrera-Martínez Y, Martín-Marcuatu JJ, Mohigefer Barrera J, Jiménez-Hoyuela JM, García Jiménez R. Hybrid image (SPECT/CT) in the early diagnosis of a patient with calciphylaxis. *Nefrologia*. 2019.
2. Sahu KK, Lal A, Mishra AK, Chastain I. Calciphylaxis: challenges in the diagnosis and management. *J Family Med Prim Care*. 2019;8:4044-5.
3. Sahu KK, Mishra AK, Lal A, Javadi H. Calciphylaxis: continues to surprise. *Nephrology*. 2020, <http://dx.doi.org/10.1111/nep.13643>.
4. Sherif AA, Sahu KK, Li Y, Javadi H. Penile Calciphylaxis: A Rare Case of Penile Lesion. *Indian J Nephrol*. 2020;30:138-140.
5. Nigwekar SU. Calciphylaxis. *Curr Opin Nephrol Hypertens*. 2017;26:276-81.

Kamal Kant Sahu^{a,*}, Amos Lal^b, Ajay Kumar Mishra^a

^a Department of Internal medicine, Saint Vincent Hospital, Worcester, Massachusetts, United States of America

^b Division of Pulmonary and Critical Care Medicine, Mayo Clinic, Rochester

* Corresponding author.

E-mail address: drkksahu85@gmail.com (K.K. Sahu).

0211-6995/© 2021 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.nefro.2020.11.019>