

errata

In the article "Assessment of the new CKD-EPI equation to estimate the glomerular filtration rate" in Revista Nefrología volume 30, issue 2, 2010, errors were discovered in Table 1 (page 187) and Table 4 (page 190). The corrections appear below:

Table 1. Estimated glomerular filtration rate formula
CKD-EPI

Black subjects

Women

- Creatinine \leq 62: estimated GFR = $166 \times ([\text{creatinine}/88.4/0.7]^{-0.329}) \times 0.993^{\text{age}}$
- Creatinine $>$ 62: estimated GFR = $166 \times ([\text{creatinine}/88.4/0.7]^{-1.209}) \times 0.993^{\text{age}}$

Men

- Creatinine \leq 80: estimated GFR = $163 \times ([\text{creatinine}/88.4/0.9]^{-0.411}) \times 0.993^{\text{age}}$
- Creatinine $>$ 80: estimated GFR = $163 \times ([\text{creatinine}/88.4/0.9]^{-1.209}) \times 0.993^{\text{age}}$

White/other subjects

Women


- Creatinine \leq 62: estimated GFR = $144 \times ([\text{creatinine}/88.4/0.7]^{-0.329}) \times 0.993^{\text{age}}$
- Creatinine $>$ 62: estimated GFR = $144 \times ([\text{creatinine}/88.4/0.7]^{-1.209}) \times 0.993^{\text{age}}$

Men

- Creatinine \leq 80: estimated GFR = $141 \times ([\text{creatinine}/88.4/0.9]^{-0.411}) \times 0.993^{\text{age}}$
- Creatinine $>$ 80: estimated GFR = $141 \times ([\text{creatinine}/88.4/0.9]^{-1.209}) \times 0.993^{\text{age}}$

GFR: glomerular filtration rate expressed in ml/min/1.73m²; creatinine expressed in µmol/l; age expressed in years.

Continued on next page>>



errata

Table 4. Concordances (shown in boldface black) in the classification of stages of chronic kidney disease (CKD) for glomerular filtration rate (GFR) estimated using the MDRD-IDMS and CKD-EPI formulas, taking GFR by MDRD-IDMS as a reference and considering sex and age variables simultaneously. In blue, subjects who were reclassified in stages with higher GFRs using the CKD-EPI formula. In normal font, subjects who were reclassified in stages with a lower GFR using the CKD-EPI formula.

			GFR (MDRD-IDMS)						
			ERC 1	ERC 2	ERC 3A	ERC 3B	ERC 4	ERC 5	
Women	<70 years	GFR (CKD-EPI)	ERC 1	572 (100)	201 (33.4)				
			ERC 2		400 (66.6)	828 (34.4)			
			ERC 3A			1,581 (65.6)	599 (24.7)		
			ERC 3B				1,824 (75.3)	300 (12.7)	
			ERC 4					2,068 (87.3)	
	>70 years		ERC 5				2 (0.1)	107 (4.1)	
			ERC 1	34 (52.3)				2,496 (95.9)	
			ERC 2	31 (47.7)	238 (100)	87 (6.5)			
			ERC 3A			1,197 (89.9)	39 (2.0)		
			ERC 3B			47 (3.5)	1,781 (92.2)	7 (0.3)	
Men	<70 years	GFR (CKD-EPI)	ERC 4				98 (5.1)	1,971 (93.9)	
			ERC 5				121 (5.8)	1,680 (100)	
			ERC 1	732 (99.3)	194 (18.8)				
			ERC 2	5 (0.7)	840 (81.2)	1,197 (24.6)			
			ERC 3A			3,669 (75.4)	678 (15.7)		
	>70 years		ERC 3B				3,624 (83.9)	230 (5.7)	
			ERC 4				16 (0.4)	3,775 (92.9)	
			ERC 5				57 (1.4)	53 (1.2)	
			ERC 1	96 (42.7)				4,346 (98.8)	
			ERC 2	129 (57.3)	637 (100)	19 (0.6)			

Values expressed as total number of cases and percentage of concordance (in parentheses).

We apologise to our readership