

<sup>b</sup> Residente de Medicina Interna Universidad Pontificia Bolivariana, Medellín, Colombia

\* Autor para correspondencia.

Correos electrónicos: [lm.serna@hotmail.com](mailto:lm.serna@hotmail.com), [johnfredynieto@gmail.com](mailto:johnfredynieto@gmail.com) (L.M. Serna-Higuaita).

<http://dx.doi.org/10.1016/j.nefro.2015.07.008>

0211-6995/© 2015 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## An unusual case of peritonitis after vaginal leak in a patient on peritoneal dialysis

### Un caso infrecuente de peritonitis tras fuga vaginal en un paciente en diálisis peritoneal

Dear Editor,

Continuous ambulatory peritoneal dialysis (CAPD) is used as an alternative to hemodialysis and has very few complications which include abdominal hernia, peritonitis, processus vaginalis, pleural leakage, and those related to the catheter exit site.<sup>1</sup> It may occasionally be complicated by leakage of dialysate fluid into the abnormal sites.<sup>2,3</sup> We report a case of a woman who experienced vaginal leakage during CAPD and after peritonitis.

A 24 year old female patient was diagnosed with end stage renal disease during pregnancy. There was no other etiological factor, but hypertension. Patient was recommended 20 hours of hemodialysis, but she refused and gave birth to a living child on the 38th gestational week. She has been on CAPD for the last 3 months. Since last week she has started to have abdominal pain accompanied with nausea and fever (>38 °C). With these complaints she was referred to our policlinic and her peritoneal fluid cell count was found 4860/mm<sup>3</sup> (95% polymorphonuclear leukocytes). Cultures were obtained and she was admitted to our hospital with the diagnosis of peritonitis. Blood analysis showed leukocytes 9000/mm<sup>3</sup>, hemoglobin: 8.8 g/dl (11.5–16), C-reactive protein: 14.2 mg/dl (0–0.5), sedimentation: 135 mm/hour. *Pseudomonas aeruginosa* was detected in blood cultures. She was on empiric ceftazidime and cefazolin treatment and treatment was continued because *P. aeruginosa* was found sensitive to this treatment. Urine culture remained sterile. Patient stated that there is dialysate in her vagina. She was consulted with obstetrics and gynecology regarding any fistulas. A urine catheter was placed and it is understood that origin of dialysate was vagina. Contrast enhanced computerized tomography was done showing a vaginal fistula (Figs. 1 and 2). CAPD catheter was removed. Surgical operation was found unnecessary and hemodialysis was started. Patient was discharged after 3 weeks of antibiotic therapy.

Patients on CAPD have an increased risk of both hernia formation and dialysate leakage as an intraperitoneal pressure related complications. One of the rare peritoneal fluid leakage is through processus vaginalis<sup>3</sup> and main



Fig. 1 – Arrows shows vaginal leak.



**Fig. 2 – Arrows shows vaginal leak.**

mechanisms include leakage through the fallopian tube and then the uterus and vagina, or leakage through a fistula between the uterus and the peritoneal cavity formed after surgery.<sup>3</sup> One of the reason for fistula formation is peritoni-

tis which is often caused by skin bacteria penetration via transvisceral, transvaginal or hematogenous spread of organisms, and the CAPD catheter can become colonized by bacteria creating a biofilm that repeatedly seeds the peritoneum.<sup>4</sup>

In our case, a rare entity; vaginal fistula and also another rarely seen clinic; peritonitis secondary to vaginal fluid were present. Probably secondary to the fistula, peritonitis developed and after peritonitis, fluid from fistula tract has increased and caused vaginal leakage.

#### BIBLIOGRAFÍA

1. Henderson S, Brown E, Levy J. Safety and efficacy of percutaneous insertion of peritoneal dialysis catheters under sedation and local anaesthetic. *Nephrol Dial Transplant.* 2009;24:3499-504.
2. Tzamaloukas AH, Gibel LJ, Eisenberg B, Goldman RS, Kanig SP, Zager PG, et al. Early and late peritoneal dialysate leaks in patients on CAPD. *Adv Perit Dial.* 1990;6:64-70.
3. Leblanc M, Ouimet D, Pichette V. Dialysate leaks in peritoneal dialysis. *Semin Dial.* 2001;14:50-4.
4. Peritoneal dialysis-associated peritonitis rates and outcomes in a national cohort are not improving in the post-millennium (2000-2007). *Perit Dial Int.* 2011;31(6):639-50.

Ali Bakan, Alihan Oral\*, Osman Kostek, Sabahat Alışır Eçder, Ali Rıza Odabaş

*Istanbul Medeniyet University, Goztepe Training and Research Hospital, Department of Internal Medicine, Kadiköy, Istanbul, Turkey*

\* Corresponding author.

E-mail address: [dr.alihanoral@gmail.com](mailto:dr.alihanoral@gmail.com) (A. Oral).

<http://dx.doi.org/10.1016/j.nefro.2015.08.006>  
0211-6995/© 2015 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).