

## Basophil activation test in interstitial nephritis. Some comments

### Test de activación de basófilos en la interstitial nephritis. Algunos comentarios

Dear Editor:

The recent paper by Lara Belmar Vega and coworkers introduced the role of a CD123<sup>pos</sup>/HLADR<sup>neg</sup> basophil activation test (BAT) in a case report of interstitial nephritis, suggesting the occurrence of an interstitial nephropathy secondary to the use of omeprazole.<sup>1</sup> Omeprazole is well known from past reports to cause interstitial nephritis.<sup>2</sup> The renal oedematosus interstitium having a massive infiltration of leukocytes and eosinophils, suggested the author to assess an immune response, due to the omeprazole therapy. This case report should assess the optimal use of BAT in omeprazole hypersensitivity.<sup>3</sup> Despite the fact that the authors were not endowed with further allergic tests, such as skin prick test (SPT) or serum IgEs to assess their evidence, they moved on the clinical suspicion of an allergy-driven nephropathy caused by an hypersensitivity response to omeprazole and found a CD63 stimulation index (SI)  $\geq 4.1$  and  $\geq 10\%$  respect the basal, non activated level.<sup>1</sup> The analytical performance of a BAT based on the CD123<sup>pos</sup>/HLADR<sup>neg</sup> gating protocol allowed the authors to easily capture basophils in a flow cytometry (FC) approach and to assess cell activation by evaluating the CD63 membrane upregulation upon the activation from omeprazole-caused drug hypersensitivity. Yet, the activation is only slightly higher (i.e. 10.25%) than the indicated cut off, which is  $\leq 5\%$  and did not reach the much more encouraging level of an fMLP-mediated activation (34.9%).<sup>1</sup> Causes underlying this moderate, poor activation may be further elucidated by introducing in the test a polyclonal anti-IgE agonist, to probe the level of releasability and activation of basophils in the IgE-mediated mechanism.<sup>4</sup> Furthermore, a CD203c marker to assess also a possible involvement of non IgE-mediated reactions might be useful to better comprehend the observed reaction to omeprazole.<sup>1</sup> Noteworthy, omeprazole targets H2-receptors, so hampering the counter-regulation of basophil activation by histamine, their constitutive activation and the subsequent down-regulation of the Fc $\epsilon$ RI/IgEs complexes, causing therefore a desensitized cell and/or a CD63 exhausted pool.<sup>5-8</sup> The apparently low level of activation can be also explained by the same gating strategy used by the authors. As already reported in our labs, capturing basophils as SSC<sup>low</sup>/CD123<sup>pos</sup> cells, results in the inclusion of further non basophilic CD123<sup>pos</sup> cells.<sup>9</sup> These cells can be differentiated as CD45<sup>pos</sup> cells and/or CD203c<sup>neg</sup> leukocytes, while basophils are notoriously CD45<sup>dim</sup>/CD123<sup>bright</sup> cells.<sup>9</sup> The inclusion of more CD123<sup>pos</sup> events in the gate,

while basophils are only Cd123<sup>bright</sup> cells, underestimates the CD63%, because a possible apparent “cellular loss”, as previously reported, actually caused by a biased gating protocol.<sup>10,11</sup>

Therefore, the enthusiastic conclusion about the use of BAT in drug hypersensitivity, should be softened by introducing some bias-preventing warnings:

- basophils should be optimally captured in FC as CD45<sup>dim</sup>/CD123<sup>bright</sup>/HLA-DR<sup>neg</sup> cells, without starting in the FSC/SSC dot plot;
- the introduction of a CD203c activation marker may be particularly useful to discern non IgE from IgE-mediated responses.

In the case report described by the authors one cannot exclude the hypothesis that the effect of omeprazole on basophils may be of immune, pharmacological origin, rather than allergic. The interaction with H2 and H4-receptors may block the histamine-mediated loop to deactivate basophils, causing the ongoing release of IL-4 from these leukocytes and the activation of a Th-mediated response, an effect that can increase the production of IgGs, with consequent nephritis onset.<sup>12,13</sup>

This paper is interesting because permits to focus onto the major effect exerted by a BAT in the clinics.

---

#### Conflict of interest

The authors state they have no conflict of interest.

---

#### BIBLIOGRAFÍA

- Belmar Vega L, López Hoyos M, San Segundo Arribas D, Irure Ventura J, Fernández Fresnedo G, Ruiz San Millán JC, et al. Basophil activation test. Tool for the diagnosis of interstitial nephritis. *Nefrologia*. 2019;39:436-8.
- Yip D, Kovac S, Jardine M, Horvath J, Findlay M. Omeprazole-induced interstitial nephritis. *J Clin Gastroenterol*. 1997;25:450-2.
- Laguna JJ, Bogas G, Salas M, Mayorga C, Dionicio J, Gonzalez-Mendiola R, et al. The basophil activation test can be of value for diagnosing immediate allergic reactions to omeprazole. *J Allergy Clin Immunol Pract*. 2018;6:1628-36, e2.

4. Chirumbolo S, Vella A, Ortolani R, De Gironcoli M, Solero P, Tridente G, et al. Differential response of human basophil activation markers: a multi-parameter flow cytometry approach. *Clin Mol Allergy*. 2008;6:12.
5. Chirumbolo S. Basophil activation test in allergy: time for an update? *Int Arch Allergy Immunol*. 2012;158:99-114.
6. Chirumbolo S, Brizzi M, Ortolani R, Vella A, Bellavite P. Inhibition of CD203c membrane up-regulation in human basophils by high dilutions of histamine: a controlled replication study. *Inflamm Res*. 2009;58:755-64.
7. Mommert S, Kleiner S, Gehring M, Eiz-Vesper B, Stark H, Gutzmer R, et al. Human basophil chemotaxis and activation are regulated via the histamine H4 receptor. *Allergy*. 2016;71:1264-73.
8. Tedeschi A, Lorini M, Arquati M, Miadonna A. Regulation of histamine release from human basophil leucocytes: role of H1, H2 and H3 receptors. *Allergy*. 1991;46:626-31.
9. Chirumbolo S, Bjorklund G, Vella A. Using a CD45dim/CD123bright/HLA-DRneg phenotyping protocol to gate basophils in FC for airway allergy. CD123 does not decrease. *Adv Respir Med*. 2017;85:193-201.
10. Santos AF, Bécáres N, Stephens A, Turcanu V, Lack G. The expression of CD123 can decrease with basophil activation: implications for the gating strategy of the basophil activation test. *Clin Transl Allergy*. 2016;6:11.
11. Chirumbolo S. Major pitfalls in BAT performance may be caused by gating protocols and CD63% cut off evaluation. *Cytometry A*. 2014;85:382-5.
12. Moharana AK, Bhattacharya SK, Mediratta PK, Sharma KK. Possible role of histamine receptors in the central regulation of immune responses. *Indian J. Physiol Pharmacol*. 2000;44:153-60.
13. Jutel M, Blaser K, Akdis CA. The role of histamine in regulation of immune responses. *Chem Immunol Allergy*. 2006;91:174-87.

Salvatore Chirumbolo<sup>a,\*</sup>, Geir Bjørklund<sup>b</sup>, Antonio Vella<sup>c</sup>

<sup>a</sup> Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy

<sup>b</sup> Council for Nutritional and Environmental Medicine (CONEM), Mo i Rana, Norway

<sup>c</sup> Unit of Immunology and Cytofluorimetry-AOUI, Policlinico GB Rossi piazzale AL Scuro 10, 37134 Verona, Italy

\*Corresponding author.

E-mail address: [salvatore.chirumbolo@univr.it](mailto:salvatore.chirumbolo@univr.it) (S. Chirumbolo).

<https://doi.org/10.1016/j.nefro.2019.06.005>

0211-6995/© 2019 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Glomerulonefritis necrosante en un paciente con VIH, VHC y leishmaniasis visceral

### Necrotizing glomerulonephritis in patients with HIV, HCV and visceral leishmaniasis

Sr. Director:

La coinfección VIH/leishmaniasis visceral es frecuente y cursa de forma tórpida y recurrente. La afectación renal incluye glomerulonefritis y afectación tubular. Presentamos un caso poco frecuente.

Se trata de un varón de 46 años adicto a alcohol y drogas parenterales diagnosticado en 2006 de infección VIH estadio C y VHC genotipo 4. Inicia TARGA en 2011, tras su primera descompensación ascítica por cirrosis hepática Child C, con hipertensión portal, esplenomegalia y pancitopenia. Lo suspende voluntariamente en mayo de 2015 y lo reinicia en noviembre del 2016 (raltegravir, abacavir y lamivudina). Un mes después, con carga viral no detectable y sin restauración inmune (CD4 74/mm<sup>3</sup>), desarrolla fracaso renal agudo no oligúrico (creatinina máxima 5,7 mg/dL), proteinuria mixta de 2 g/día, microhematuria y tumoración en el dorso de la lengua (fig. 1) con biopsia de displasia epitelial severa y candidiasis mucosa. En las pruebas complementarias destacaba

descenso de C'3, gammapatía policlonal, aumento de inmunoglobulinas y de cadenas kappa y lambda y ANA positivos (1/640). Las crioglobulinas, el Mantoux y las serologías (VHB, sífilis, *Toxoplasma*) fueron negativas. Se determinaron IgG de CMV y *Leishmania* y antígeno rK39 en sangre, que resultaron positivos. La PCR para *Leishmania* en la lengua y la médula ósea fueron negativas. Se realizó biopsia renal, en la que se identificaron 6 glomérulos, uno con focos de necrosis segmentaria con fibrina y 2 con semilunas celulares sin trombos (fig. 2). En el intersticio aparecía inflamación crónica moderada y fibrosis, sin alteraciones tubulares. No había signos de vasculitis. La IFD era positiva de predominio mesangial para C'3 (++) , IgG (+), C1q e IgM (+), y negativa para IgA y cadenas ligeras. No se identificó amiloide con la técnica del Rojo Congo. ME no estaba disponible. El paciente inició tratamiento con anfotericina B liposomal, normalizando 3 semanas más tarde la función renal y mejorando las alteraciones urinarias. Dos meses después inició tratamiento