

Original article

Analysis of emergency Department Frequentation among patients with advanced CKD (chronic kidney disease): Lessons to optimize scheduled renal replacement therapy initiation[☆]

Víctor Lorenzo Sellarés

Servicio de Nefrología, Hospital Universitario de Canarias, La Laguna (Santa Cruz de Tenerife), Spain

ARTICLE INFO

Article history:

Received 3 May 2017

Accepted 9 May 2018

Available online 27 November 2018

Keywords:

Advanced chronic kidney disease

Renal replacement therapy initiation

Emergency Department frequentation

Non-scheduled renal replacement therapy initiation

ABSTRACT

The decision to initiate renal replacement therapy (RRT) implies a wide margin of uncertainty. Glomerular filtration rate (GFR) tells us the magnitude of renal damage. Proteinuria indicates the speed of progression. However, nowadays more than 50% of patients are still initiating RRT hastily, and it is life threatening.

Hypothesis: By analysing Emergency Department (ED) frequentation and causes of a hurried initiation, we can better schedule the timing of the start of RRT.

Method: Retrospective and observational study of all CKD patients in our outpatient clinic. ED frequentation and hospitalization (Hos) time were reviewed during a 12-month period. We analyzed: (1) time at risk, purpose (modality of RRT), previous comorbidity; (2) causes of ED frequentation and Hos; (3) type of initiation: “scheduled” vs. “non-scheduled”, and within these “non-planned” vs. “potentially planned”.

Results: Of a total of 267 patients (time at risk 63.987 days, 70 ± 13 years, 67% males, 38% diabetics), 68 (25%) patients came to hospital on 97 occasions: 39 only ED, 46 ED+Hos and 12 only Hos. ED frequentation was one patient every 4.3 days, and bed occupation was almost 3 per day. Main causes: 47% cardiopulmonary (1/3 heart failure), 11% vascular peripheral+cerebral, 11% gastrointestinal: 8/11 due to bleeding (all with anticoagulants/antiplatelet agents). Thirty-one (12%) patients initiated RRT: of these, 14 (45%) were scheduled (6 PD, 6 HD, and 2 living donor RTx), and 17 (55%) were not scheduled or were rushed, all with venous central catheter. Following the objectives of this study, the non-scheduled group were itemized into 2 groups: 9 non-planned (initial indication of conservative management or patient’s refusal to undergo dialysis, and diverse social circumstances not controllable by the nephrologist) and 8 were considered potentially planned (6 heart failure, one gastrointestinal bleeding and one peripheral vascular complication). This last group (potentially planned), when compared with the 14 patients who started

DOI of original article:

<https://doi.org/10.1016/j.nefro.2018.05.006>.

[☆] Please cite this article as: Lorenzo Sellarés V. Análisis de la frecuentación de Urgencias en consulta ERCA (enfermedad renal crónica avanzada): enseñanzas para optimizar el inicio programado en tratamiento renal sustitutivo. Nefrología. 2018;38:622–629.

E-mail address: vls243@gmail.com

2013-2514/© 2018 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

treatment in a scheduled manner, had significant differences in that they were older, with more previous cardiac events, and GFR almost double that of the other group. All of them started treatment in the ED.

Conclusion: This analysis provides us with knowledge on those patients who may benefit from an earlier preparation in RRT. We suggest that patients with previous cardiac events, especially with a risk of gastrointestinal bleeding, should start the preparation for RRT even with GFR rates of 20–25 ml/min. In spite of the retrospective nature of this study, and taking into account the difficulties of carrying out clinical trials in this population, we propose this suggestion as complementary to the current recommendations for a scheduled start using this technique.

© 2018 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Análisis de la frecuentación de Urgencias en consulta ERCA (enfermedad renal crónica avanzada): enseñanzas para optimizar el inicio programado en tratamiento renal sustitutivo

R E S U M E N

Palabras clave:

Enfermedad renal crónica avanzada
Inicio de tratamiento renal sustitutivo
Frecuentación de Urgencias
Inicio no programado en tratamiento renal sustitutivo

La decisión de empezar tratamiento renal sustitutivo (TRS) conlleva un amplio margen de incertidumbre. El filtrado glomerular (FG) nos dice la magnitud del daño. La proteinuria, la velocidad de progresión. A pesar de estas premisas, más del 50% de los pacientes continúan iniciando TRS de forma precipitada y con riesgo vital.

Hipótesis: Analizando la frecuentación de Urgencias (Urg) y las causas determinantes de un inicio precipitado, podremos programar mejor el momento de iniciar un TRS.

Método: Estudio retrospectivo, observacional, de la frecuentación de Urg y del tiempo de hospitalización (Hos) de todos los pacientes de la consulta ERCA, durante un período de 12 meses. Se analizó: 1) tiempo en riesgo, destino (modalidad de TRS), comorbilidad previa. 2) Causas de frecuentación de Urg y Hos. 3) Tipo de inicio: «programado» vs. «no programado» y, dentro de estos, «no planificables» vs. «potencialmente planificables».

Resultados: De 267 pacientes (con un tiempo en riesgo de 63.987 días; 70 ± 13 años; 67% varones; 38% diabéticos), 68 (25%) pacientes acudieron al hospital en 97 ocasiones: 39 solo Urg, 46 Urg + Hos y 12 solo Hos. La frecuentación de Urg fue de un paciente cada 4,3 días y la ocupación de camas fue de casi 3 diarias. Causas predominantes: 47% cardiopulmonar (1/3 insuficiencia cardíaca), 11% vascular periférico + cerebral, 11% digestivo: 8/11 por sangrado (todos con anticoagulantes/antiagregantes). Iniciaron TRS: 31 (12%): de estos, 14 (45%) de forma programada (6 DP, 6 HD y 2 TxR de donante vivo); 17 (55%) no programados o precipitados, todos con catéter venoso. Siguiendo los objetivos del estudio, estos últimos se desglosaron en 2 grupos: 9 no planificables (indicación inicial de manejo conservador o negativa del paciente a dializarse, y circunstancias sociales diversas no controlables por el nefrólogo) y 8 que consideramos potencialmente planificables (6 con fallo cardíaco, uno con hemorragia digestiva y uno vascular periférico). Estos últimos (potencialmente planificables), comparados con los 14 que iniciaron de forma programada, tenían significativamente mayor edad, más eventos cardíacos previos y el FG casi duplicaba al del otro grupo; todos entraron por Urg.

Conclusión: Este análisis nos aporta conocimiento sobre aquellos pacientes que pueden beneficiarse de una preparación más precoz en TRS: proponemos que en los enfermos con eventos cardíacos previos, especialmente con riesgo de sangrado digestivo, se inicie la preparación para TRS aun con tasas de FG de 20-25 ml/min. A pesar de la naturaleza retrospectiva del estudio y ante la dificultad práctica de ensayos clínicos en esta población, proponemos esta medida como complemento a las recomendaciones actuales para un inicio programado en esta técnica.

© 2018 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

It is well known that early programme for the initiation of renal replacement therapy (RRT) has a favourable impact on the patient's survival.^{1,2} However, the decision to start RRT implies a wide margin of uncertainty on these patients with a precarious health. It should be kept in mind that starting early may deprive the patient from a period of a time with a good quality of dialysis free. The fact is that it is being debated that early onset improves later survival.³⁻⁵ Therefore, the responsible doctor usually faces the difficult balance between starting too early, or too late; requiring a central venous catheter, often with vital risk.

The KDIGO Guidelines⁶ emphasize the initiation of RRT in symptomatic patients and/or in the presence of a glomerular filtration rate (GFR) of 5–10 mL/min. In general, these situations result from late referral to the nephrologist.^{2,7-9} But this will not be our scenario; now will concentrate in patients being followed in of advanced chronic kidney disease outpatient clinics (ACKD). Usually these patients manifest a subjective feeling of being well, maintain a relatively high urinary output, and are especially reticent to be dialysed. However, frequently an adverse event forces an emergency hemodialysis through a catheter. The KDIGO Guidelines⁶ present an outline for the risk of end-stage renal disease. However, despite these premises, the proportion of patients who start precipitously and with a venous catheter is very high, around 50%, in most series.¹⁰⁻¹³ In addition, the DOPPS study reports unscheduled initiation rates with venous catheter between 23 and 70%.¹⁴

Our hypothesis is that the analysis of emergencies room visits and the causes that determine a precipitous onset of dialysis may help to optimize the plan to start RRT in a scheduled manner.

The objective of the present study is to analyze retrospectively, the frequency of emergency room visits and their relationship with the precipitated onset in RRT, in patients from our ACKD clinic (monocentric study) during a period of 12 months.

Method

The study was performed in a ACKD outpatient clinic of a reference Hospital. The centre covers the health care of approximately 400,000 people. Clinical Records are available from 1994 to the end of 2015 from a total of 1,492 patients. The modalities of RRT offered to the patients are hemodialysis (HD), peritoneal dialysis (PD) and kidney and pancreas transplantation (Tx) from living or cadaveric donor. Inhabitants of this region are characterized for a high incidence of early onset of diabetes which has led to a high prevalence of diabetic nephropathy (DN) associated with high cardiovascular comorbidity.^{1,15-17}

The ACKD clinics attend patients with GFR <30 mL/min, and other candidates to receive RRT. Most patients come from other outpatient clinics, or after a hospital discharge of cases with potentially irreversible CKD. Patients, who return to dialysis after the loss of the kidney graft, are previously treated in the transplant clinic and they are not included in this study.




Fig. 1 – Patients at risk and outcomes in the clinic of advanced chronic kidney disease (ACKD) during 2015.

PCD: primary care doctor. LFU: loss of follow-up. RRT: renal replacement therapy. HD: hemodialysis. PD: peritoneal dialysis.

TxR: kidney transplant. Time at risk: period of time in days, when patients are in ACKD consultation during 2015.

The patients visit the emergency room or are admitted to the same hospital where patients are being followed as outpatient.

The standard protocols recommended by the Clinical Practice Guidelines are implemented in our clinic. Likewise, we consider important to instruct patients on the adjustment of drugs (diuretics, antihypertensives) both in case of hydrosaline retention and also prevent volume depletion (summer season, gastroenteritis, etc.).^{1,18} To optimize tracking and adherence to our recommendations, we recommend visits on demand and routine visits which scheduled bimonthly and quarterly and a telephone number is provided to the patient.

Design

Retrospective, observational and single centre study of the frequency of emergency room attendance and time of hospitalization of all patients of the ACKD outpatient clinic (monocentric study) during a period of 12 months. Fig. 1 shows the number of patients attended in consultation during 2015, the period of time at risk and the outcome during this period. As compared with previous years (only as a reference), we found that the average age of patients was increasing, with more male predominance, and the high rate of ND was maintained as the primary cause of renal disease among incident patients^{15,16} (Table 1).

The diagnoses of comorbidity and relevant biochemical parameters were extracted from the patients electronic records. A cardiologist followed up all the patients with previous cardiovascular comorbidity. The diagnoses of congestive heart failure (CHF) was based on clinical data and a previous echocardiography. Also, to assess cardiac function the cardiologist recommended stress or pharmacological tests, and catheterization. The coincidence of several factors was highlighted, often coexisting the diagnosis of arrhythmia, a variable degree systolic or diastolic dysfunction, underlying coronary disease; pulmonary hypertension, and the

Table 1 – General demographic data, cardiovascular comorbidity and relevant biochemical parameters, grouped according to whether they had been in ER or admitted to the Hospital during 2015.

	All 267	Not-admitted 199 (75%)	Admitted 68 (25%)	
Age	70 ± 13	69 ± 14	73 ± 11	0.007
Gender (%male)	67	66	68	NS
No. diabetics (%)	38	35	48	0.045
Cardiovascular comorbidity (%)	51	41	81	0.001
Coronary disease	50	50	51	
Valvulopathy	14	16	12	
CHF	42	37	49	
Albumin (g/dL)	4.2 ± 0.4	4.3 ± 0.3	3.9 ± 0.4	0.01
Haemoglobin (g/dL)	11.8 ± 1.3	12.11 ± 1.2	11.1 ± 1.4	0.01
HbA1c (%)	7.0 ± 1.5	7.1 ± 1.6	7.0 ± 1.1	NS

CHF: congestive heart failure.

aggravating effect of previous pulmonary disease (history of smoking, with diagnosis or suspicion of chronic obstructive pulmonary disease). Likewise, the accompanying gastrointestinal pathology (angiodyplasia, diverticula, polyps, etc.), was a clinical concern due to the frequent use of anticoagulants and anaemia due to digestive haemorrhage.

Initiation of RRT

Patients were divided into 3 groups: (1) Programmed: they started with access to dialysis or received early TxR, in a planned manner. (2) Not planned: those patients who due to special, social, health or clinical circumstances, were outside the control of a nephrologist, and started dialysis using venous catheter in a precipitated manner. These circumstances are specified in the Results. (3) Abrupt and/or urgent initiation due to clinical events, however that could have benefited from an advance programming in RRT. This is the group of special interest in the analysis of the frequency of emergency room visits.

Emergency Room visits

Emergency frequentation and/or hospitalization was obtained from the electronic records of the hospital.

All patients had a clinical history at admission and clinical evolution, with analytical data and imaging studies. Information collected included the reason to attend the emergency room (ER), the clinical services involved in the care of the patient, the evolution and the final outcome of the patient. The patients were initially assessed by the emergency doctor, and thereafter by the nephrologist and any other specialist.

Statistics

Given the nature of the study, does not involve a statistical analysis. Only some parameters are described as mean and standard deviation.

Results

In Fig. 1 it is highlighted that 31 patients (12%) initiated RRT; and 11 (4%) died. Of these 11 deaths, five occurred during the




Fig. 2 – Frequency of visits to the emergency room and hospitalizations: number of patients and time of bed occupation.

emergency assistance, four due to cerebral vascular event and one was of cardiac origin.

Frequentation of Emergency Room and Hospitalization

In Table 1 it is shown the basic data of patients at risk during 2015; patients were grouped into those admitted to the hospital at least on one occasion, and those who were not admitted during that period. Sixty eight out of the 267 patients went to the Emergency Room or were hospitalized (25%). These 68 patients visited the hospital on 97 occasions: 50 once, 10 twice, 6 three times, one four and one five times. The patients who were hospitalized were adult, with high proportion of DN and high cardiovascular comorbidity. The values of haemoglobin and albumin were significantly lower in those admitted, but within a range close to normal. The control of diabetes based on HbA1c levels was not different between groups.

Fig. 2 shows that from the 97 hospital visits, 85 (87%) were to the ER, and 46 (47%) were hospitalized. Twelve patients were hospitalized without going to ER; none of them required RRT during hospitalization.

Table 2 – Causes of emergency attendance and Hospitalization grouped by relevant pathology and frequency.

	n	%
Cardiac	46	47.4
Cerebrovascular	6	6.2
Peripherovascular	5	5.2
Gastrointestinal	11	11.3
Nephrology	9	9.3
Others	20	20.6
Total	97	100.0

The visits to ER of this population was one every 4.3 days, which is 1.6% of the time at risk of the patients in medical consultation. Likewise, there were almost 3 hospital beds occupied daily.

The causes of ER frequentation and Hospitalization are described in Table 2. Almost half of the patients had an event of cardiac or cardiopulmonary origin, dyspnoea was the predominant manifestation (67%) (frequently referring to "shortness of breath"), orthopnoea, most cases were grouped under the heading of congestive heart failure (CHF), pulmonary oedema or respiratory failure. All had been included within the so-called cardiorenal syndrome.¹⁹ Added to 11 admissions that were due to peripheral or cerebral vascular causes, the hospitalizations of cardiovascular origin reached almost 58%.

Another important cause of admission was gastrointestinal (GI) disturbance, which accounted 11% of the hospitalizations. It is remarkable that 8 were for some type of GI bleeding, all undergoing anticoagulation or antiaggregant therapy. Half of these patients had symptoms of heart failure. Also, there were two other hospitalizations precipitated by cerebral vascular bleeding who were also on anticoagulants.

Nine patients were admitted in the Nephrology service with deterioration of renal function, of which six started dialysis. Some patients were also admitted in "others" speciality wards due to various pathologies and none of them required dialysis.

There was a high percent of patients on anticoagulant/antiaggregant medication; among not hospitalized patients, 65% received anticoagulation/antiaggregant therapy and among hospitalized patients a 78% were on anticoagulation/antiaggregant therapy. The proportion of those receiving acenocoumarol (Sintrom®) was significantly increased in hospitalized patients (24% vs 11%, $p < 0.007$).

Initiation of renal replacement therapy

Renal replacement therapy (RRT) was initiated in 31 patients (Fig. 3). In Fourteen (45%) patients the type of RRT had been planned and had arteriovenous fistula (AVF) or a functioning peritoneal catheter or an anticipated Kidney transplant. However, two of them started HD after visiting ER due to CHF. Nine of the patients who had a precipitous initiation of hemodialysis with a venous catheter, were on circumstances that can be considered virtually unavoidable.

The second group is the most important for our analysis, since we consider that they were susceptible to earlier programming, thus avoiding the implant of a catheter. In these patients the initiation of dialysis was precipitated by cardiovascular causes.

Comparison of these two groups is shown in Table 3, the 14 patients starting dialysis in a programmed manner with AVF, peritoneal catheter in place or anticipated renal transplant. There were 8 additional patients that had to start regular hemodialysis using tunnelled catheter; they were seen in the ER and it is important to keep in mind that 80% had a his-




Fig. 3 – Proportion of patients who started RRT, scheduled vs. unscheduled or precipitated initiation. Within this later group, patients were grouped in “unable to programme RRT” and “programmable RRT”. PC: peritoneal catheter. CHF: congestive heart failure. TC: tunnelled catheter ER: start from the Emergency Room.

Table 3 – Comparison of patients that started on a scheduled basis versus those that we consider not programmed, but potentially programmable.

	Programmed	Programmable	p
n	14	8	
Age	57 ± 10	77 ± 10	0.0001
Start in the emergency department	2 (15%)	8 (100%)	0.0001
Previous cardiovascular disease	2 (14%)	7 (88%)	0.0001
Pre initial MDRD	8.9 ± 2.5	14.8 ± 4.8	0.001

CV: cardiovascular; MDRD: Modification of Diet in Renal Disease Study equation.³⁶

tory of previous cardiac events and three patients had 2 or 3 hospitalizations due to the same cause during the last year. However, the GFR measured in the outpatient clinic was twice as high as those who initiated dialysis on scheduled basis; thus, such a rapid progression of the kidney damage was not expected.

Discussion

We know that the GFR and proteinuria are the best markers of kidney damage and progression of disease.^{20–25} The GFR reflects magnitude of damage, and proteinuria is associated to the rapidity of progression.^{26–28} However, although its predictive value is unquestionable, they do not prevent a hasty start in a considerable proportion of patients.

A revision of the literature reveals that despite efforts to achieve a programmed initiation of RRT, more than half of the patients start RRT precipitously through a central venous catheter.^{10–13} The objective proposed by guidelines and documents is that 80% of the incidents in dialysis should start dialysis with a definitive access; however this objective appears to be difficult,²⁹ although it is a priority in the management of this population. The increased age and increasing comorbidity, especially cardiovascular comorbidity, may not allow to achieve this objective despite the efforts by the professionals. In fact, heart failure is the main cause of hospitalization in patients 65 years and older, with a high rate of readmissions.³⁰ The recent study by Ronksley PE and col³¹ on renal patients confirm that 52% of emergency frequentations were due to CHF episodes.

There is abundant information about the importance of early referral to the nephrologist, so the initiation of dialysis is organized.^{2,8–10} But also, transcendent circumstances have been verified, beyond the moment of deriving the patient.^{12,32} A study similar to ours, by Gomis-Couto and col,¹² analyzed the causes of not programmed initiation of dialysis, although they did not address the emergency room attendance. These authors emphasize that unexpected incidents and the accelerated aggravation of the ERC as the most non-preventable variable. But very important, the authors note that the greatest area for improvement lies in the delay in performing vascular access. In another study, Mendelssohn DC and col³² highlighted that 37% of patients started dialysis while hospitalized, and 54% with a temporary catheter; and that this is independent of early or late referral. In this regard, Bhan V and col³³ emphasize that the delay in performing vascular access in fast progressors is an important preventable obstacle for a programmed initiation of dialysis.

The period of preparation to start RRT is variable among the centres, but not less than the two months required for maturation of AVF or for peritoneal catheter ready to use or even more time in the anticipated TxR.

The analysis of the frequency of ER visits and the reasons for precipitated initiation of dialysis may provide useful information. By reviewing the literature, we found that the issue of early initiation of dialysis was not approached from this perspective. The present study has provided information about the use of Hospital Emergency Services by patients with ACKD. Out of a population at risk of 267 patients, the frequency of ER visits was one every 4 days approximately, and the occupation of hospital beds was almost 3 beds/day. We did not find similar information from other sources, but the annual analysis should allow to analyze annual changes, and evaluate the results after improving strategies.

Our study also shows that despite the high age and comorbidity, the proportion of patients who died or who started RRT appear to be low (Fig. 1), however we have no other data to compare. This observation may be similar – or better – in other health areas, but in any case, what indicates is that progress in health care entail a greater survival, a greater use of resources, and very importantly, generates an extraordinary health care expense that must adapt to these growing demand.^{34,35}

The next step was to analyze the circumstances that precipitated the unscheduled initiation of dialysis and make an attempt to advance in the preparation of these patients. As also observed by Gomis-Couto A and col,¹² there are situations as frequent as unavoidable described in Fig. 3 that precipitates the need for dialysis. The most characteristic cases are those patients that state “I will not be dialyzed” an idea that changes as soon as they find themselves in the ER symptomatic and at risk of losing their life, sudden renal deterioration due to an unpredictable intercurrent process; and failure or delay in preparation for access for dialysis. Other potential causes described in the study by Gomis-Couto A,¹² such as acute renal failure that do not recover renal function or late referral to the nephrologist, have not occurred, since our practice receives patients that have already passed that filter.

The most interesting part of our study was the search of those cases that could have benefited from an early planning for dialysis. There is a sum of “extrarenal” factors which have been decisive for the hasty and not programmed beginning of dialysis; cardiovascular events accounted for six out of ten visits to the ER with precipitated start of hemodialysis with catheter (Table 2). This should warn the nephrologist

of the need to speed up the preparation of patients with this clinical profile; despite the frequent subjective sensation of well-being between events, and with levels of renal function “acceptable”. In these cases, the most characteristic were the repeated episodes of CHF, the result of a frequent combination of underlying coronary disease, valvular disease and heart rhythm disorders; that frequently coexisting with chronic lung disease and gastrointestinal pathology. The last one is very important, since the frequent prescription of anticoagulants or antiaggregants, increases the risk of gastrointestinal bleeding, anaemia and heart failure.

The blood parameters (haemoglobin and serum albumin) were slightly lower in those patients that frequently visited the ER, but within a range close to normal, hence being helpful in warning about these events. Fig. 4 shows a diagram of the most frequent comorbidity profile that leads the ACKD patient to go to the Emergency Service symptomatic and initiate dialysis in a precipitated and unscheduled manner.

Coming back to our hypothesis, this analysis of the frequency of ER visits provides additional information to the glomerular filtration and proteinuria as the main variables of renal risk. As a corollary, we suggest that in circumstances illustrated in Fig. 4, especially if they have led the patient to attend the emergency department repeatedly in the last year, preparation for RRT should begin even with glomerular filtration rates of 20–25 mL/min. Very especially for those patients with resistant proteinuria of medium-high range (more than 1.5 g/24 h and very especially if it is persistently greater than 2.5 g). The early preparation of these cases does not imply initiating dialysis immediately, but giving priority to the preparation of the chosen modality.

Despite the retrospective nature of the study, and given the practical difficulty of preparing and obtaining results from prospective studies, we propose this opinion as a complement to the current recommendations for a scheduled initiation in technique.




Fig. 4 – Diagram of the most frequent comorbidity profile that leads the patient to the emergency room symptomatic and requiring initiation of RRT in a hasty way. ACS: Acute Coronary Syndrome. Dysf: dysfunction (S: systolic; D: diastolic). CHF: congestive Heart Failure.

Conflicts of interest

The author has no conflicts of interest to declare.

REFERENCES

- Lorenzo V. [Chronic renal failure outpatient clinic. A 12 years experience]. *Nefrologia*. 2007;27:425–33 [in Spanish].
- Lorenzo V, Martn M, Rufino M, Hernandez D, Torres A, Ayus JC. Predialysis nephrologic care and a functioning arteriovenous fistula at entry are associated with better survival in incident hemodialysis patients: an observational cohort study. *Am J Kidney Dis*. 2004;43:999–1007.
- Korevaar JC, Jansen MA, Dekker FW, Jager KJ, Boeschoten EW, Krediet RT, et al. Netherlands Cooperative Study on the Adequacy of Dialysis Study G: When to initiate dialysis: effect of proposed US guidelines on survival. *Lancet*. 2001;358:1046–50.
- Korevaar JC, Jansen MA, Dekker FW, Boeschoten EW, Bossuyt PM, Krediet RT, et al. Evaluation of DOQI guidelines: early start of dialysis treatment is not associated with better health-related quality of life. *Am J Kidney Dis*. 2002;39:108–15.
- Traynor JP, Simpson K, Geddes CC, Deighan CJ, Fox JG. Early initiation of dialysis fails to prolong survival in patients with end-stage renal failure. *J Am Soc Nephrol*. 2002;13:2125–32.
- KDIGO Clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl*. 2013;3:1–150.
- Gorriz JL, Sancho A, Pallardo LM, Amoedo ML, Barril G, Salgueira M. Longer pre-dialysis nephrological care is associated with improved long-term survival of dialysis patients. More facts. *Nephrol Dial Transplant*. 2002;17:1354–5.
- Stack AG. Impact of timing of nephrology referral and pre-ESRD care on mortality risk among new ESRD patients in the United States. *Am J Kidney Dis*. 2003;41:310–8.
- Winkelmayer WC, Glynn RJ, Levin R, Owen WF Jr, Avorn J. Determinants of delayed nephrologist referral in patients with chronic kidney disease. *Am J Kidney Dis*. 2001;38:1178–84.
- Gorriz JL, Sancho A, Pallardo LM, Amoedo ML, Martin M, Sanz P, et al. [Prognostic significance of programmed dialysis in patients who initiate renal substitutive treatment. Multicenter study in Spain]. *Nefrologia*. 2002;22:49–59 [in Spanish].
- Castellano I, Gallego S, Labrador PJ, Gomez-Martino JR, Covarsi A. [The start of renal replacement therapy in a Spanish department]. *Nefrologia*. 2006;26:445–51 [in Spanish].
- Gomis Couto A, Teruel Briones JL, Fernandez Lucas M, Rivera Gorin M, Rodriguez Mendiola N, Jimenez Alvaro S, et al. Causes of unplanned hemodialysis initiation. *Nefrologia*. 2011;31:733–7.
- Rodriguez JA, Lopez J, Cleries M, Vela E. Vascular access for haemodialysis: an epidemiological study of the Catalan Renal Registry. *Nephrol Dial Transplant*. 1999;14:1651–7.
- Ethier J, Mendelssohn DC, Elder SJ, Hasegawa T, Akizawa T, Akiba T, et al. Vascular access use and outcomes: an international perspective from the Dialysis Outcomes and Practice Patterns Study. *Nephrol Dial Transplant*. 2008;23:3219–26.
- Lorenzo V, Boronat M. [Terminal kidney disease associated with diabetes in the Canary Islands: a public health problem with high economic cost and human suffering]. *Nefrologia*. 2010;30:381–4 [in Spanish].

16. Lorenzo V, Boronat M, Saavedra P, Rufino M, Maceira B, Novoa FJ, et al. Disproportionately high incidence of diabetes-related end-stage renal disease in the Canary Islands. An analysis based on estimated population at risk. *Nephrol Dial Transplant*. 2010;25:2283-8.
17. Lorenzo V, Sanchez E, Vega N, Hernandez D. Renal replacement therapy in the Canary Islands: demographic and survival analysis. *J Nephrol*. 2006;19:97-103.
18. Lorenzo V. Doctor, how much should I drink? *Nefrologia*. 2014;34:693-7.
19. Ronco C, McCullough PA, Anker SD, Anand I, Aspromonte N, Bagshaw SM, et al. Cardiorenal syndromes: an executive summary from the consensus conference of the Acute Dialysis Quality Initiative (ADQI). *Contrib Nephrol*. 2010;165:54-67.
20. Astor BC, Matsushita K, Gansevoort RT, van der Velde M, Woodward M, Levey AS, et al. Lower estimated glomerular filtration rate and higher albuminuria are associated with mortality and end-stage renal disease. A collaborative meta-analysis of kidney disease population cohorts. *Kidney Int*. 2011;79:1331-40.
21. Brenner BM. Retarding the progression of renal disease. *Kidney Int*. 2003;64:370-8.
22. Gansevoort RT, Matsushita K, van der Velde M, Astor BC, Woodward M, Levey AS, et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes. A collaborative meta-analysis of general and high-risk population cohorts. *Kidney Int*. 2011;80:93-104.
23. Palmer BF. Proteinuria as a therapeutic target in patients with chronic kidney disease. *Am J Nephrol*. 2007;27:287-93.
24. Ruggenti P, Perna A, Mosconi L, Pisoni R, Remuzzi G. Urinary protein excretion rate is the best independent predictor of ESRF in non-diabetic proteinuric chronic nephropathies. *Gruppo Italiano di Studi Epidemiologici in Nefrologia (GISEN)*. *Kidney Int*. 1998;53:1209-16.
25. Ruggenti P, Perna A, Remuzzi G. Retarding progression of chronic renal disease: the neglected issue of residual proteinuria. *Kidney Int*. 2003;63:2254-61.
26. Palmer BF. Supratherapeutic doses of angiotensin receptor blockers to decrease proteinuria in patients with chronic kidney disease. *Am J Nephrol*. 2008;28:381-90.
27. Parving HH, Hovind P, Rossing K, Andersen S. Evolving strategies for renoprotection: diabetic nephropathy. *Curr Opin Nephrol Hypertens*. 2001;10:515-22.
28. Ruggenti P, Schieppati A, Remuzzi G. Progression, remission, regression of chronic renal diseases. *Lancet*. 2001;357:1601-8.
29. Lopez Revuelta K, Barril G, Caramelo C, Delgado R, Garcia Lopez F, Garcia Valdecasas J, et al. [Developing a clinical performance measures system for hemodialysis, Quality Group Spanish Society of Nephrology]. *Nefrologia*. 2007;27:542-59 [in Spanish].
30. Grancelli H, Varini S, Ferrante D, Schwartzman R, Zambrano C, Soifer S, et al. Randomized trial of telephone intervention in chronic heart failure (DIAL): study design and preliminary observations. *J Card Fail*. 2003;9:172-9.
31. Ronksley PE, Tonelli M, Manns BJ, Weaver RG, Thomas CM, MacRae JM, et al. Emergency department use among patients with CKD: a population-based analysis. *Clin J Am Soc Nephrol*. 2017;12:304-14.
32. Mendelssohn DC, Curtis B, Yeates K, Langlois S, MacRae JM, Semeniuk LM, et al. Suboptimal initiation of dialysis with and without early referral to a nephrologist. *Nephrol Dial Transplant*. 2011;26:2959-65.
33. Bhan V, Soroka S, Constantine C, Kiberd BA. Barriers to access before initiation of hemodialysis: a single-center review. *Hemodial Int*. 2007;11:349-53.
34. Lorenzo V, Perestelo L, Barroso M, Torres A, Nazco J. [Economic evaluation of haemodialysis. Analysis of cost components based on patient-specific data]. *Nefrologia*. 2010;30:403-12 [in Spanish].
35. Lorenzo-Sellares V, Pedrosa MI, Santana-Exposito B, Garcia-Gonzalez Z, Barroso-Montesinos M. Cost analysis and sociocultural profile of kidney patients. Impact of the treatment method. *Nefrologia*. 2014;34:458-68.
36. Levey AS, Coresh J, Greene T, Stevens LA, Zhang YL, Hendriksen S, et al., Chronic Kidney Disease Epidemiology Collaboration. Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med*. 2006;145:247-54.